



## Psychobiology of Pain and Exercise in Gulf War Veterans with Chronic Musculoskeletal Pain

**Dane B. Cook**  
William S. Middleton Memorial Veterans  
Hospital, Madison, WI  
University of Wisconsin - Madison



## Chronic musculoskeletal pain in Gulf War Veterans

- 15% (100,000 of ~700,000) report chronic muscle pain symptoms (Kang et al., 2000)
- One of three major factors of Gulf War illness (Fukuda et al., 1997).
- Reported twice as frequently (OR=3.06) in Gulf War Veterans (GVs) than non-GVs (Kang et al., 2000; Thomas et al., 2006)
- Estimated 1 in 7 seek health care for war-related concerns and 12% receive disability compensation (Engel et al., 2004; DVA report, 1998; Hodgson & Kipen, 1999; Kang et al., 2000)
- Follow-up data indicate that symptoms have not resolved (Blanchard et al., 2006; Ozakinci et al., 2006; Thomas et al., 2006)



## Today's Presentation

- Exercise alters pain sensitivity in Gulf War Veterans (GVs) with chronic musculoskeletal pain (CMP)
  - Follow-up functional brain imaging study
- Functional brain imaging of chronic musculoskeletal pain
  - Past, present and future
- Overview of Merit Review exercise training project
  - Exercise, pain, brain function & structure

## Exercise alters pain sensitivity in Gulf War Veterans with chronic muscle pain

*Journal of Pain, 2010*

*Supported by Department of Veteran Affairs Grant #  
561-00215*

### Pathophysiological mechanisms that may maintain CMP

- Fibromyalgia (FM):
  - Chronic musculoskeletal pain disorder
- The mechanism for maintenance of pain and other symptoms in FM is unknown
- Research suggests that FM pain is maintained by abnormal central nervous system regulation of sensory stimuli
  - Behavioral data
    - More sensitive to experimental pain stimuli
    - Altered modulation – DNIC & wind-up
    - **Do not exhibit exercise-induced hypoalgesia (EIH)**



## Purpose & Hypotheses

- To determine the impact of an acute bout of exercise on pain sensitivity in GVs with CMP compared to healthy GVs
- **GVs with CMP:**
  - H1: Report lower pain thresholds and higher pain ratings than healthy GVs
  - H2: Rate naturally occurring muscle pain during exercise as more intense than healthy GVs
  - H3: Not demonstrate EIH, but instead become more sensitive to experimental pain stimuli following acute exercise

## Methods

- **N = 32 participants (WRIISC)**
  - n= 15 GVs with widespread & chronic muscle pain
  - n= 17 GVs healthy and without pain
- **Testing to occur on 2 separate days**
  - Maximal exercise testing (ACSM)
  - Submaximal exercise testing and pain psychophysics
    - Psychophysical pain assessment
    - Exercise @ 70% of peak oxygen consumption for 30 minutes followed by 3-minute active recovery
    - Psychophysical pain assessment





# Results

**Table 1. Demographic, Exercise and Clinical Characteristics for the Final Sample (N = 27)**

|   | CMP (N = 11) | HEALTHY (N = 16) | E.S. |
|---|--------------|------------------|------|
| Age (y)   | 39.4 (±7.4)  | 40.9 (±7.9)      | -.20 |
| Height (cm)   | 176.2 (±8.6) | 174.6 (±9.2)     | .19  |
| Weight (kg)   | 96.5 (±25.8) | 90.8 (±14.2)     | .30  |
| Resting HR (bpm)  | 72 (±13)     | 65 (±11)         | .63  |
| Resting SBP (mmHg)  | 117 (±10)    | 116 (±9)         | .12  |
| Resting DBP (mmHg)  | 75 (±7)      | 77 (±7)          | -.37 |
| VO <sub>2peak</sub> (mL·kg <sup>-1</sup> ·min <sup>-1</sup> ) | 28.2 (±7.7)  | 31.9 (±8.5)      | -.47 |
| Peak PO (watts)   | 169.6 (±45)  | 204.0 (±53)      | -.69 |
| Average PO (Submax)   | 82.8 (±31)   | 111.7 (±33)      | -.90 |
| Widespread pain complaints                                    | 11/11        | 0/16             | —    |
| Pain in 4 body quadrants                                      | 9/11         | 0/16             | —    |
| FM diagnosis  | 3/11         | 0/16             | —    |
| Current pain intensity (0–5)                                  | 3.27 (±1.1)  | —                | —    |



Abbreviations: CMP, Chronic muscle-pain patients; Healthy, Healthy controls; E.S., Effect size (calculated as Cohen's  $d = \frac{\bar{x}_1 - \bar{x}_2}{s_{pooled}}$ ); HR (bpm), Heart rate (beats per minute); SBP, Systolic blood pressure; DBP, Diastolic blood pressure; VO<sub>2peak</sub>, Volume of O<sub>2</sub> consumption at peak effort; PO, Power output (Watts attained at peak during maximal exercise and average watts during 30-minute sub-max exercise test).  
 NOTES: Current pain intensity represents pain felt in muscles and joints at clinical intake, 0 = no pain, 1 = mild pain, 2 = moderate pain, 3 = substantial pain, 4 = severe pain, and 5 = very severe.  
 Values in columns headed with groups represent means (±standard deviation) or proportions.

**Table 2. Comparison of Pre- and Post-Exercise Heat and Pressure-Pain Thresholds Across Groups**


|                         | CMP (N = 11) | HEALTHY (N = 16) | E.S. |
|-------------------------|--------------|------------------|------|
| Heat-Pain Threshold     |              |                  |      |
| Pre-exercise (°C)       | 42.9 (±3.3)  | 44.1 (±3.3)      | -.36 |
| Post-exercise (°C)      | 43.4 (±2.8)  | 44.8 (±2.8)      | -.52 |
| Pressure-Pain Threshold |              |                  |      |
| Pre-exercise (sec)      | 22.9 (±16.0) | 46.9 (±41.1)     | -.75 |
| Post-exercise (sec)     | 31.5 (±38.3) | 49.1 (±38.6)     | -.48 |

Abbreviations: CMP, Chronic muscle-pain patients; Healthy, Healthy controls; E.S., Effect size (calculated as Cohen's  $d = \frac{\bar{x}_1 - \bar{x}_2}{s_{pooled}}$ ).  
 NOTES: Values in columns headed with group names represent means (±standard deviation) for respective variables. The means in this table were compared with a Group × Trial (2 × 2) repeated-measures ANOVA with no significant main effects or interactions found. Effect sizes for the within-group differences across trials (pre- vs post-exercise) were all small (ie,  $d < .3$ ).


## GVs with CMP perceive exercise as more painful and effortful than healthy GV's



### GVs w/ CMP are more sensitive to heat pain than healthy GV's and become more sensitive following acute exercise



### GVs with CMP demonstrated large increases in affective pain ratings from pre- to post-exercise



## Conclusions

- GVs with CMP:
  - Are more sensitive to experimental heat pain stimuli than healthy GVs
  - Perceive sub-maximal exercise as more painful & effortful than healthy GVs
  - Describe experimental pain stimuli as more intense and more unpleasant following 30 minutes of moderately intense, submaximal exercise


## Take Home Point

- Data are consistent with psychophysical & exercise literature for FM & suggest that the central nervous system of GVs with CMP are not properly regulating sensory information.
- GVs with CMP do not exhibit EIH, but instead become hyperalgesic following an acute bout of exercise.



# Functional Imaging of Pain in Veterans with Unexplained Muscle Pain

*Department of Veteran Affairs Grant:  
Merit Review Entry Program Project*



## Take Home Point

- Similar to our work in FM, GVs with CMP exhibit augmented brain responses to both non-painful and painful sensory stimuli.
- It is currently unclear whether this is a result of enhanced processing or decreased regulation of nociceptive information.

## Imaging the cognitive modulation of pain in CMP

*Supported by:*  
*Department of Veteran Affairs Grant # 561-00436*  
*&*  
*NIH (NIAMS) RO1 AR050969*

## Determine the influence of anticipation & attention on brain responses to pain

- Anticipation manipulated by randomly assigning participants to 'pain' and 'no pain' conditions
- Attention manipulated by having participants complete the Stroop color-word task while receiving painful stimuli

## Results: Anticipation & Stroop

**COGNITIVE MODULATION OF PAIN INTENSITY**

$p < 0.05$

accounting for Stroop alone and pain alone

**CO**

Thalamus, ACC, DLPFC, Insula


**FM**

Thalamus, VLPFC, PAG

FIBROMYALGIA (N=17)

HEALTHY CONTROL (N=18)

**Diffusion tensor imaging data demonstrating decreased fractional anisotropy and increased mean diffusivity in chronic muscle pain**




## Take Home Point

- It appears that patients with CMP are less efficient at regulating pain.
- This may be in part due to poor communication between brain regions involved in descending pain control.
- Augmented sensory processing and inefficient regulation may be one mechanism through which CMP may be maintained.

# The impact of resistance exercise training on pain and brain function in GV's with CMP

Supported by:  
 Department of Veteran Affairs Merit Review Award

## Mechanistic Resistance Exercise Training Trial



# Funding Acknowledgement



**Thank You**