

INPATIENT MEDICATIONS

PHARMACIST'S USER MANUAL

Version 5.0 January 2005

(Revised January 2012)

Revision History

Each time this manual is updated, the Title Page lists the new revised date and this page describes the changes. If the Revised Pages column lists "All," replace the existing manual with the reissued manual. If the Revised Pages column lists individual entries (e.g., 25, 32), either update the existing manual with the Change Pages Document or print the entire new manual.

Date	Revised	Patch	Description
	Pages	Number	
01/2012	i, v-vii, 10,	PSJ*5*254	Updated Table of Contents Added Order Checks/Interventions (OCI) to "Hidden Actions" section
	21 25		Defined OCI Indicator Updated Schedule Type text
	29		Updated text under Interventions Menu
	42a, 49, 56, 56a, 75, 89, 99, 106-		Updated Pharmacy Interventions for Edit, Renew, and Finish orders for Unit dose and IV
	106b,		
	124c		Added note to Drug-Drug Interactions
	124f-124g		Added note to Drug-Allergy Interactions
	124k-124l 124x		Updated Allergy/ADR Example Order Checks Added "Display Pharmacist Intervention" section
	124x 124y-124z		Defined Historical Overrides/Interventions
	224, 228,		Updated Glossary
	232, 233,		
	234		
	239-244		Updated Index REDACTED
09/2011	58	PSJ*5*235	Updated 'Note' section regarding Expected First Dose REDACTED
07/2011	i, 16	PSJ*5*243	Update Revision History Update Index
	246		Revised the existing display in the <i>Non-Verified/Pending</i>
			Orders [PSJU VBW] option from a pure alphabetic listing of
			patient names, to a categorized listing by priority. Added
			"priority" to Index.
04/2011		DCI+5+101	REDACTED
04/2011	i	PSJ*5*181	Updated Revision History
	-		
	17		Updated: Example: Patient Information Screen
	19		Update: "Select DRUG"
			Note was updated
	21		
	27.29		
			Update: "Select DRUG"

Date	Revised Pages	Patch Number	Description
	31		Updated: Example: Edit an Intervention
	32		Updated: Example: Delete an Intervention
	33		Updated: Example: View an Intervention
	34		Updated: Example: Print an Intervention
	35-36b		New: Discontinued Codes and Example of Inpatient Order
			Entry
	37		New: Example: Patient Information
	40		Updated: 4.1.5.1 Discontinue
	41		Updated: Example: Discontinue an Order (continued)
	46		Updated: Example: Verify an Order (continued)
	61-62b		Updated: 4.1.8 Inpatient Profile, Discontinued Codes, &
			example
	65		Updated: Example: Patient Information
	66		Updated: Example: Patient Record
	67		Updated: Example: Patient Information
	73-74		Updated: Example: New Order Entry
	76		Updated: Example: New Intervention
	77		Updated: Example: Edit an Intervention
	78		Updated: Example: Delete an Intervention
	79		Updated: Example: View an Intervention
	80		Updated: Example: Print an Intervention
	81		Updated: 4.2.3.5 View Profile
	83		Updated: Example: Patient Information
	98		Updated: 4.1.5.4 Hold
	118		Updated text
	120-120b		Updated: 4.2.7. Inpatient Profile, Discontinued Codes, &
			example
	122		Updated: Example: Inpatient Profile
	123-124v		Updated: 4.3. Order Checks
	125		Added Note
	136		Updated: Example: Extra Units Dispensed Report
	137		Updated: Example: Reporting Medication Returns
	153		Updated: Example: Patient Profile
	190		Updated: Example: Extended Patient Profile Report
	192a-192b		Updated: 8.1.5. Patients on Specific Drug(s)
	194-195		Updated: Example: IV Individual Labels
	196-196d		New: Example: IV Individual Labels (Print New Labels)
	219-220		New: 10. CPRS Order Checks – How They Work
	221-222		New: 11. Error Messages
	223-238		Updated: Glossary page numbering
	239-246		Updated: Index & page numbering
			REDACTED REDACTED
9/2010	i-ii, 174	PSJ*5*232	Deleted paragraph referring to Start/Stop date prompts of
			Action Profile #1 option as this is not how the option works.
			REDACTED
			THE PROTEIN

Date	Revised Pages	Patch Number	Description
06/2010	i-v, 33-34, 25a-25d, 124a-124b, 124e-124f, 239-241	PSJ*5*113	Added new Order Validation Requirements. Removed Duplicate Order Check Enhancement functionality, (removed in a prior patch). REDACTED
02/2010	i-ii, iv-v, 192a-b, 214a-b, 239-241	PSJ*5*214	Updated Table of Contents to include new sections. Added new sections 8.1.5 and 8.2.4 to reference <i>Patients on Specific Drug(s)</i> option that is now commonly used by pharmacists who may have been assigned this option directly and not as part of the Supervisor's Menu. Added <i>Patients on Specific Drug(s)</i> option to the Index. REDACTED
12/2009	56, 56a, 56b iii	PSJ*5*222	Added description of warning displayed when finishing a Complex Unit Dose Order with overlapping admin times. Corrected page numbers in Table of Contents. REDACTED
07/2009	43	PSJ*5*215	When Dispense Drug is edited for an active Unit Dose, an entry is added to the activity log. REDACTED
02/2009	226	PSJ*5*196	Update to IV Duration REDACTED
0829 /2008	iii, 20-27, 54, 68-76, 94-95, 104- 106, 236, 240-241	PSJ*5*134	Inpatient Medication Route changes added, plus details on IV type changes for infusion orders from CPRS, pending renewal functions, and expected first dose changes. REDACTED
07/2007	iii, 124 a-d 5, 17-18, 27-28, 30-34, 37- 38, 65-68, 76-80, 83-84, 119- 120, 123- 124, 149- 150, 195- 196, 209-210	PSJ*5*175 PSJ*5*160 PSJ*5*145	Modified outpatient header text for display of duplicate orders. Added new functionality to Duplicate Drug and Duplicate Class Order Check definitions. Modifications for remote allergies, to ensure all allergies are included when doing order checks using VA Drug Class; Analgesic order checks match against specific class only; check for remote data interoperability performed when entering patient's chart; and list of remote allergies added to Patient Information screen. REDACTED
07/2007	155a-155b, 162a-162b, 168a-168b	rsj*3*145	On 24-Hour, 7-Day, and 14-Day MAR Reports, added prompt to include Clinic Orders when printing by Ward or Ward Group. Also added prompt to include Ward Orders when printing by Clinic or Clinic Group. REDACTED

Date	Revised Pages	Patch Number	Description
05/2007	25	PSJ*5*120	Modified Inpatient Medications V. 5.0 to consider the duration the same way as all other stop date parameters, rather than as an override. REDACTED
12/2005	1, 124-124b	PSJ*5*146	Remote Data Interoperability (RDI) Project: Removed document revision dates in Section 1. Introduction. Updated Section 4.3. Order Checks to include new functionality for checking allergies, drug reactions, and interactions. REDACTED
03/2005	iv-vii, 114-116, 223, 236-241	PSJ*5*112	Updated TOC to correct Index page number. (p. iv) In Unit Dose Menu Tree, changed Clinic Stop Dates to Clinic Definition. (p. v) In Section 1., Introduction, updated revision dates and added reference to Release Notes. (p. 1) In Sections 4.2.5.1., 4.2.5.3., and 4.2.5.3., added a sentence that refers to the IMO parameter NUMBER OF DAYS UNTIL STOP from the CLINIC DEFINITION file. (p.114-116) Updated Glossary; added definition for CLINIC DEFINITION File. (p. 223) Updated Index; added CLINIC DEFINITION file and Inpatient Medication Orders for Outpatients page number references; reflowed all following Index pages. (p. 236-241) REDACTED
01/2005	All	PSJ*5*111	Reissued entire document to include updates for Inpatient Medication Orders for Outpatients and Non-Standard Schedules. REDACTED

Table of Contents

1.	Introduction	1	
2.	Orientation	3	
3.	List Manager		
	3.1. Using List Manager	7	
	3.2. Hidden Actions	7	
4.	Order Options	11	
	4.1. Unit Dose Medications Option	11	
	4.1.1. Order Entry	12	
	4.1.2. Non-Verified/Pending Orders		
	4.1.3. Inpatient Order Entry	17	
	4.1.4. Patient Actions	18	
	4.1.5. Order Actions	40	
	4.1.6. Discontinue All of a Patient's Orders	60	
	4.1.7. Hold All of a Patient's Orders	60	
	4.1.8. Inpatient Profile	61	
	4.2. IV Menu Option	63	
	4.2.1. Order Entry (IV)		
	4.2.2. Inpatient Order Entry	65	
	4.2.3. Patient Actions	66	
	4.2.4. Order Actions	86	
	4.2.5. IV Types	114	
	4.2.6. Profile (IV)	117	
	4.2.7. Inpatient Profile	120a	
	4.3. Order Checks	123	
	4.3.1. Inpatient Duplicate Therapy	124h	
	4.3.2 Discontinuing Duplicate Inpatient Orders	i	
	4.3.4. Sample Drug/Drug Interactions	124p	
	4.3.5. Sample Therapeutic Order Check Displays	124u	
	4.3.6. Display of Provider Overrides and Pharmacist Interventions	X	
5.	Maintenance Options	125	
	5.1. Unit Dose	125	
	5.1.1. Edit Inpatient User Parameters	125	
	5.1.2. Edit Patient's Default Stop Date	126	

	5.2. IV	126
	5.2.1. Change Report/Label Devices (IV)	126
	5.2.2. Change to Another IV Room (IV)	127
6.	PIck List Menu	129
	6.1. PIck List	129
	6.2. ENter Units Dispensed	
	6.3. EXtra Units Dispensed	
	6.4. Report Returns	
	6.5. Reprint Pick List	138
	6.6. Send Pick List To ATC	140
	6.7. Update Pick List	141
7.	Production Options	143
	7.1. Ward List (IV)	143
	7.2. Update Daily Ward List (IV)	144
	7.3. Manufacturing List (IV)	146
	7.4. RETurns and Destroyed Entry (IV)	148
	7.5. Barcode ID – Return and Destroy (IV)	152
8.	Output Options	153
	8.1. Unit Dose	
	8.1.1. PAtient Profile (Unit Dose)	
	8.1.2. Reports Menu	
	8.1.3. Align Labels (Unit Dose)	
	8.1.4. Label Print/Reprint	
	8.1.5. Patients on Specific Drug(s)	
	8.2. IV	
	8.2.1. Label Menu (IV)	
	8.2.2. REPorts (IV)	
	8.2.3. SUSpense Functions (IV)	208
9.	Inquiries Option	215
	9.1. Unit Dose	
	9.1.1. INQuiries Menu	215
	9.2. IV	
	9.2.1. Drug Inquiry (IV)	217

10.	CPRS Order Checks – How They Work	219
	10.1. Order Check Data Caching	219
11.	Error Messages	221
	11.1. Error Information	222
12.	Glossary	223
13.	Index	239

Synonym	Action	Description
IN	Intervention Menu	Displays, allows actions to be taken on
		orders where interventions are required or suggested.
PRO	Patient Profiles	Displays the Patient Profile Menu
IP	Inpatient Medications Profile	Generates an Inpatient Profile for a patient
IV	IV Medications Profile	Generates an IV Profile for a patient
UD	Unit Dose Medications Profile	Generates a Unit Dose Profile for a patient
OP	Outpatient Prescriptions	Generates an Outpatient Profile for a patient
AP1	Action Profile #1	Generates an Action Profile #1
AP2	Action Profile #2	Generates an Action Profile #2
EX	Patient Profile (Extended)	Generates an Extended Patient Profile
CWAD	CWAD Information	Displays the crises, warnings, allergies, and
		directives information on a patient

The Intervention menu hidden action is available to the Medication Profile and Detailed Order List Manager screens when utilizing the following options:

- *Inpatient Order Entry* [PSJ OE]
- Non-Verified/Pending Orders [PSJU VBW]
- Order Entry [PSJU NE]
- Order Entry (IV) [PSJI ORDER]

The following actions are available while in the Unit Dose Order Entry Profile.

Synonym	<u>Action</u>	Description
DC	Speed Discontinue	Speed discontinue one or more orders
		(This is also available in the <i>Inpatient</i>
		Order Entry and Order Entry (IV) options.)
RN	Speed Renew	Speed renewal of one or more orders
SF	Speed Finish	Speed finish one or more orders
SV	Speed Verify	Speed verify one or more orders

The following actions are available while viewing an order.

Synonym	<u>Action</u>	Description
CO	Copy an order	Allows the user to copy an active,
		discontinued, or expired Unit Dose order
DIN	Drug Restriction/Guideline	Displays the Drug Restriction/Guideline
	Information	Information for both the Orderable Item and
		Dispense Drug
I	Mark Incomplete	Allows the user to mark a Non-Verified
		Pending order incomplete
JP	Jump to a Patient	Allows the user to begin processing another
		patient
N	Mark Not to be Given	Allows the user to mark a discontinued or
		expired order as not to be given

<u>Synonym</u>	<u>Action</u>	<u>Description</u>
OCI	Order Checks/Interventions	Indicates there are associated CPRS Overrides and/or Pharmacist Interventions. When the OCI indicator displays on the Order Detail screen, the user can type "OCI" to display associated CPRS Provider Overrides and/or Pharmacist Interventions.

If the Dispense Drug or Orderable Item has a non-formulary status, this status will be displayed on the screen as "*N/F*" beside the Dispense Drug or Orderable Item.

Order Checks/Interventions (OCI) Indicator:

When the OCI indicator displays on the Order Detail screen, it indicates there are associated CPRS Provider Overrides and/or Pharmacist Interventions for this order. The Order Checks/Interventions indicator **OCI**> will display on the same line as the Orderable Item field, to the left of the drug text indicator **DIN**> (if it exists).

```
*(1)Orderable Item: METRONIDAZOLE TAB
                                                                  <OCI><DIN>
     Instructions: 250MG
*(2)Dosage Ordered: 250MG
                              (3)Start: 07/11/11 15:33
REQUESTED START: 07/11/11 16:00
         Duration:
*(4) Med Route: ORAL
                                            (5) Stop: 07/25/11 15:33
 (6) Schedule Type: CONTINUOUS
*(8) Schedule: Q36H
(9) Admin Times:
* (10)
      Provider: PSJPROVIDER, ONE [es]
 (11) Special Instructions:
 (12) Dispense Drug
                                                  U/D
                                                           Inactive Date
     METRONIDAZOLE 250MG TAB
                                                  1
     Enter ?? for more actions
        Enter ?? for more actions

AC ACCEPT
ED Edit
Select Item(s): Next Screen// AC ACCEPT
```

If the OCI indicator displays on the Order Detail screen, the user can type "OCI" to display the current CPRS Provider Overrides and/or Pharmacist Interventions associated with the order, as well as any historical overrides and interventions, if applicable.

• "DOSAGE ORDERED:" (Regular and Abbreviated)

To allow pharmacy greater control over the order display shown for Unit Dose orders on profiles, labels, MARs, etc., the DOSAGE ORDERED field is not required if only one Dispense Drug exists in the order. If more than one Dispense Drug exists for the order, then this field is required.

When a Dispense Drug is selected, the selection list/default will be displayed based on the Possible Dosages and Local Possible Dosages.

Example: Dispense Drug with Possible Dosages

```
Select DRUG: BACLOFEN 10MG TABS MS200
...OK? Yes// <Enter> (Yes)

Now Processing Enhanced Order Checks! Please wait...

Press Return to continue...

Available Dosage(s)

1. 10MG
2. 20MG

Select from list of Available Dosages or Enter Free Text Dose: 1 10MG

You entered 10MG is this correct? Yes// <Enter>
```

All Local Possible Dosages will be displayed within the selection list/default.

Example: Dispense Drug with Local Possible Dosages

```
Select DRUG: GENTAMICIN CREAM 15GM DE101 DERM CLINIC ONLY
...OK? Yes// <Enter> (Yes)

Now Processing Enhanced Order Checks! Please wait...

Press Return to continue...

Available Dosage(s)

1. SMALL AMOUNT
2. THIN FILM

Select from list of Available Dosages or Enter Free Text Dose: 2 THIN FILM

You entered THIN FILM is this correct? Yes// <Enter>
```

Note: If an order contains multiple Dispense Drugs, Dosage Ordered should contain the total dosage of the medication to be administered.

• "SCHEDULE TYPE:" (Regular)

This defines the type of schedule to be used when administering the order. If the Schedule Type entered is One-time, the ward parameter, DAYS UNTIL STOP FOR ONE-TIME, is accessed to determine the stop date. When the ward parameter is not available, the system parameter, DAYS UNTIL STOP FOR ONE-TIME, will be used to determine the stop date. When neither parameter has been set, one-time orders will use the ward parameter, DAYS UNTIL STOP DATE/TIME, to determine the stop date instead of the start and stop date being equal. When a new order is entered or an order entered through CPRS is finished by pharmacy, the default Schedule Type is determined as described below:

- If no Schedule Type has been found and a Schedule Type is defined for the selected Orderable Item, that Schedule Type is used for the order.
- If no Schedule Type has been found and the schedule contains PRN, the Schedule Type is PRN.
- Schedules meant to cause orders to display as ON CALL in BCMA must be defined in the ADMINISTRATION SCHEDULE (#51.1) file with a schedule type equal to "ON CALL."
- For all others, the Schedule Type is CONTINUOUS.



Note: During backdoor order entry, the Schedule Type entered is used unless the schedule is considered a ONE-TIME schedule. If so, the Schedule Type is changed to ONE TIME.

• "ADMINISTRATION TIME:" (Regular)

This defines the time(s) of day the order is to be given. Administration times must be entered in a two or four digit format. If multiple administration times are needed, they must be separated by a dash (e.g., 09-13 or 0900-1300). This field must be left blank for odd schedules, e.g., Q16H. If the schedule for the order contains "PRN", all Administration Times for the order will be ignored. In new order entry, the default Administration Times are determined as described below:

- If Administration Times are defined for the selected Orderable Item, they will be shown as the default for the order.
- If Administration Times are defined in the INPATIENT WARD PARAMETERS file for the patient's ward and the order's schedule, they will be shown as the default for the order.
- If Administration Times are defined for the Schedule, they will be shown as the default for the order.

• Order Validation Checks:

The following order validation checks will apply to Unit Dose orders and to intermittent IV orders.

Note: IV orders do not have Schedule Type.

Order Validation Check One

For intermittent IV orders, references to an order's Schedule Type will refer to either the TYPE OF SCHEDULE from the Administration Schedule file (#51.1), or PRN for schedule names in PRN format, or CONTINUOUS for schedule names in Day of Week format.

Order Validation Check Two

The system shall use the schedule type of the schedule from the Administration Schedule file independent of the schedule name when processing an order to determine if administration times are required for a particular order.

• Order Validation Check Three

If an order has the Schedule Type of Continuous, the Schedule entered is NOT in Day of Week (Ex. MO-FR) or PRN (Ex. TID PC PRN) format, and the frequency associated with the schedule is one day (1440 minutes) or less, the system will not allow the number of administration times associated with the order to be greater than the number of administration times calculated for that frequency. The system will allow for the number of administration times to be LESS than the calculated administration times for that frequency but not less than one administration time. (For example, an order with a schedule of BID is associated with a frequency of 720 minutes. The frequency is divided into 1440 minutes (24 hours) and the resulting calculated administration time is two. For this order, the number of administration times allowed may be no greater than two, but no less than one. Similarly, a schedule frequency of 360 minutes must have at least one administration time but cannot exceed four administration times.)

If an order has the Schedule Type of Continuous, the Schedule entered is NOT in Day of Week (Ex. MO-FR) or PRN (Ex. TID PC PRN) format, and the frequency associated with the schedule is **greater than one day** (1440 minutes) and evenly divisible by 1440, only one administration time is permitted. (For example, an order with a schedule frequency of 2880 minutes must have ONLY one administration time. If the frequency is greater than 1440 minutes and not evenly divisible by 1440, no administration times will be permitted.)

The system shall present warning/error messages to the user if the number of administration times is less than or greater than the maximum admin times calculated for the schedule or if no administration times are entered. If the number of administration times entered is less than the maximum admin times calculated for the schedule, the warning message: "The number of admin times entered is fewer than indicated by the schedule." shall appear. In this case, the user will be allowed to continue after the warning. If the number of administration times entered is greater than the maximum admin times calculated for the schedule, the error message: "The number of admin times entered is greater than indicated by the schedule." shall appear. In this case, the user will not be allowed to continue after the warning. If no admin times are entered, the error message: "This order requires at least one administration time." shall appear. The user will not be allowed to accept the order until at least one admin time is entered.

Example: New Order Entry (continued)

```
Pre-Exchange DOSES: <Enter>
ORDER VERIFIED.

Enter RETURN to continue or '^' to exit:
```

4.1.4.3. Detailed Allergy/ADR List

The Detailed Allergy/ADR List action displays a detailed listing of the selected item from the patient's Allergy/ADR List. Entry to the *Edit Allergy/ADR Data* option is provided with this list also.

Enter/Edit Allergy/ADR Data

Provides access to the Adverse Reaction Tracking (ART) package to allow entry and/or edit of allergy adverse reaction data for the patient. See the Allergy package documentation for more information on Allergy/ADR processing.

Select Allergy

Allows the user to view a specific allergy.

4.1.4.4. Intervention Menu

This option is only available to those users who hold the PSJ RPHARM key.

The Intervention Menu action allows entry of new interventions and existing interventions to be edited, deleted, viewed, or printed. Each kind of intervention will be discussed and an example will follow.

Note: Interventions can also be dynamically created in response to Order Checks for critical drug-drug interactions and allergy/ADRs. Refer to Section 4.3 Order Checks.

If a change is made to an intervention associated to an inpatient order made in response to critical drug-drug and/or allergy/ADR, the changes are reflected and displayed whenever interventions display.

New interventions entered via the Intervention Menu are at the patient level and are not associated with a particular order. Consequently, new entries made through this menu are not reflected in the OCI listing, the BCMA Display Order detail report, and do not cause highlighting in BCMA.

New: This option is used to add an entry into the APSP INTERVENTION file.

Example: New Intervention

```
Patient Information Sep 22, 2000 08:03:07 Page: 1 of 1
                               Ward: 1 West
PSJPATIENT2, TWO
   | JPATIENT2, TWO | Ward: 1 West | PID: 000-00-0002 | Room-Bed: A-6 |
                                                       Ht(cm): 167.64 (04/21/99)
   DOB: 02/22/42 (58)
                                                      Wt(kg): 85.00 (04/21/99)
   Sex: MALE
                                                    Admitted: 09/16/99
                                           Last transferred: ****
   Dx: TEST PATIENT
Allergies - Verified: CARAMEL, CN900, LOMEFLOXACIN, PENTAMIDINE, PENTAZOCINE,
                         CHOCOLATE, NUTS, STRAWBERRIES, DUST
         Non-Verified: AMOXICILLIN, AMPICILLIN, TAPE, FISH,
                         FLUPHENAZINE DECANOATE
               Remote:
   Adverse Reactions:
 Inpatient Narrative: Inpatient narrative
Outpatient Narrative: This is the Outpatient Narrative. This patient doesn't
like waiting at the pickup window. He gets very angry.
Enter ?? for more actions
PU Patient Record Update
PU Patient Record Update NO New Order Entry DA Detailed Allergy/ADR List IN Intervention Menu
                                          NO New Order Entry
VP View Profile
Select Action: View Profile// IN Intervention Menu
                             --- Intervention Menu ---
    Delete Pharmacy Intervention
Edit Pharmacy Intervention
Enter Pharmacy Intervention
                                                Print Pharmacy Intervention
DI
                                          PO
                                          VP View Pharmacy Intervention
ED
NE Enter Pharmacy Intervention
Select Item(s): NE Enter Pharmacy Intervention
Select APSP INTERVENTION INTERVENTION DATE: T SEP 22, 2000
  Are you adding 'SEP 22, 2000' as a new APSP INTERVENTION (the 155TH)? No// Y
  (Yes)
   APSP INTERVENTION PATIENT: PSJPATIENT2, TWO
                                                  02-22-42 000000002
SC VETERAN
   APSP INTERVENTION DRUG: WAR
    1 WARFARIN 10MG BL100
2 WARFARIN 10MG U/D BL100
3 WARFARIN 2.5MG BL100
4 WARFARIN 2.5MG U/D BL100
5 WARFARIN 2MG BL100TAB
s <RETURN> to see more.
                                                 TAB
                                                   TAB **AUTO STOP 2D**
                                    BL100
                                                   TAB
                                       BL100
                                                       TAB **AUTO STOP 2D**
Press <RETURN> to see more, '^' to exit this list, OR
CHOOSE 1-5: 1 WARFARIN 10MG BL100
PROVIDER: PSJPROVIDER, ONE
                                   PROV
INSTITUTED BY: PHARMACY// <Enter> PHARMACY
INTERVENTION: ALLERGY RECOMMENDATION: NO CHA
                   NO CHANGE
WAS PROVIDER CONTACTED: N NO
RECOMMENDATION ACCEPTED: Y YES
FINANCIAL COST:
REASON FOR INTERVENTION:
 1>
ACTION TAKEN:
 1>
CLINICAL IMPACT:
 1>
FINANCIAL IMPACT:
Select Item(s):
```

Note: The first time a field marked with an asterisk (*) is selected for editing, if CPRS Provider Overrides and/or Pharmacist Interventions exist for the order, entering Y (Yes) at the prompt: "Order Check Overrides/Interventions exist for this order. Display? (Y/N)? Y//" displays the following

Heading information first, followed by a summary of the Current CPRS Order Checks overridden by the Provider, as well as the Overriding Provider, plus title, Override Entered By, plus title, Date/Time Entered, and the Override Reason.

Example: Edit an Order with Provider Overrides/Interventions

```
______
      ** Current Provider Overrides for this order **
Overriding Provider: PSJPROVIDER, ONE (PROVIDER)
Override Entered By: PSJPROVIDER, ONE (PROVIDER)
 Date/Time Entered: 07/11/11 09:45
   Override Reason: testing functionality of PO & PI
 CRITICAL drug-drug interaction: TAMOXIFEN CITRATE 10MG TAB and WARFARIN NA
  (GOLDEN STATE) 1MG TAB [ACTIVE] - The concurrent use of tamoxifen or
  toremifene may increase the effects of anticoagulants. - Monograph Available
 SIGNIFICANT drug-drug interaction: TAMOXIFEN CITRATE 10MG TAB and
 THIORIDAZINE HCL 10MG TAB [UNRELEASED] - Concurrent use of inhibitors of CYP
 P-450-2D6 may decrease the effectiveness of tamoxifen in preventing breast
 cancer recurrence. Concurrent use of amiodarone or thioridazine may increase
 the risk of potentially life-threatening cardiac arrhythmias, including
 torsades de pointes. - Monograph Available
Press RETURN to Continue or '^' to Exit:
      ** Current Pharmacist Interventions for this order **
Intervention Date/Time: 07/11/11 09:50
Pharmacist: PSJPHARMACIST, ONE
                                          Drug: TAMOXIFEN CITRATE 10MG TAB
Instituted By: PHARMACY
Intervention: CRITICAL DRUG INTERACTION
Originating Package: INPATIENT
```

4.1.5.5. Renew

Medication orders (referred to in this section as orders) that may be renewed include the following:

- All non-complex active Unit Dose and IV orders.
- Orders that have been discontinued due to ward transfer or treating specialty change.
- Expired orders containing an administration schedule (Unit Dose and scheduled IV orders) that have not had a scheduled administration time since the last BCMA action was taken.
- Expired orders not containing an administration schedule (continuous IV orders) that
 have had an expired status less than the time limit defined in the EXPIRED IV TIME
 LIMIT field in the PHARMACY SYSTEM file.

Note: Complex Orders may only be renewed if all associated child orders are renewable.

Renewing Orders with CPRS Overrides/Pharmacist Interventions

When renewing an order, if CPRS Provider Overrides and/or Pharmacy Interventions exist for the order, entering Y (Yes) at the prompt: "Order Check Overrides/Interventions exist for this order. Display? (Y/N)? Y//" displays the heading information first, followed by a summary of the Current CPRS Order Checks overridden by the Provider.

If current Pharmacist Interventions exist, they will display with the following fields (if populated), Heading, Intervention Date/Time, Provider, Pharmacist, Drug, Instituted By, Intervention, Recommendation, and Originating Package.

Example: Renew an Order with Provider Overrides/Interventions

Note: When Renewing an Order in Inpatient Medications, if Current CPRS Provider Overrides do not exist and Pharmacist Interventions do exist for the order, the following displays:

```
** CURRENT PROVIDER OVERRIDES FOR THIS ORDER **

NO PROVIDER OVERRIDES TO DISPLAY
```

```
** Current Pharmacist Interventions for this order **

Intervention Date: 07/11/11 14:55

Provider: PSJPROVIDER, ONE Pharmacist: PSJPHARMACIST, ONE

Drug: WARFARIN NA (GOLDEN STATE) 1MG TAB
Instituted By: PHARMACY
Intervention: CRITICAL DRUG INTERACTION
Recommendation: OTHER Originating Package: INPATIENT
Other For Recommendation:
TEST INTERVENTION FOR CRITICAL DRUG-DRUG
```

Renewing Active Orders

The following applies when the RN (Renew) action is taken on any order with a status of "Active":

- A new Default Stop Date/Time is calculated for the order using the same calculation applied to new orders. The starting point of the Default Stop Date/Time calculation is the date and time that the order was signed in CPRS or the date and time that the RN (Renew) action was taken in Inpatient Medications.
- The RN (Renew) action does not create a new order.
- The Start Date/Time is not available for editing when an order is renewed.

Note: Orders having a schedule type of One-Time or On Call must have a status of "Active" in order to be renewed.

Renewing Discontinued Orders

IV and Unit Dose orders that have been discontinued, either through the (DC) Discontinue action or discontinued due to edit, cannot be renewed.

IV and Unit Dose medication orders that have been discontinued due to ward transfer or treating specialty change will allow the (RN) Renew action.

Renewing Expired Unit Dose Orders

The following applies to expired Unit Dose orders having a schedule type of Continuous or PRN.

- 1. The RN (Renew) action will not be available on an order with a status of "Expired" if either of the following two conditions exist:
 - a. If the difference between the current system date and time and the last scheduled administration time is greater than the frequency of the schedule. This logic will be used for schedules with standard intervals (for example, Q7H).
 - b. If the current system date and time is greater than the time that the next dose is due. This logic is used for schedules with non-standard intervals (for example, Q6H 0600-1200-1800-2400).
- 2. A new Default Stop Date/Time is calculated for the order using the same calculation applied to new orders. The starting point of the Default Stop Date/Time calculation is the date and time that the order was signed in CPRS or the date and time that the RN (Renew) action was taken in Inpatient Medications.
- 3. The (RN) Renew action does not create a new order.
- 4. The Start Date/Time is not available for editing when an order is renewed.
- 5. The renewed order has a status of "Active."

4.1.5.6. Activity Log

This action allows viewing of a long or short activity log, dispense log, or a history log of the order. A short activity log only shows actions taken on orders and does not include field changes. The long activity log shows actions taken on orders and does include the requested Start and Stop Date/Time values. If a history log is selected, it will find the first order, linked to the order where the history log was invoked from, then show an order view of each order associated with it, in the order that they were created. When a dispense log is selected, it shows the dispensing information for the order.

Example: Activity Log

```
ACTIVE UNIT DOSE
                Sep 21, 2000 12:44 25
                                                           Page:
PSJPATIENT1, ONE
                              Ward: 1 EAST
   PID: 000-00-0001
                            Room-Bed: B-12
                                                 Ht (cm):
   DOB: 08/18/20 (80)
                                                 Wt(kg):
*(1)Orderable Item: AMPICILLIN CAP
      Instructions:
 *(2) Dosage Ordered: 500MG
                                                    *(3)Start: 09/07/00 15:00
          Duration:
 *(4) Med Route: ORAL
                                                   *(5) Stop: 09/21/00 24:00
  (6) Schedule Type: CONTINUOUS
 *(8) Schedule: QID
 (9)
      Admin Times: 01-09-15-20
* (10)
      Provider: PSJPROVIDER, ONE [es]
 (11) Special Instructions:
 (12) Dispense Drug
                                                   U/D
                                                              Inactive Date
AMPICILLIN 500MG CAP
                                                   1
         Enter ?? for more actions
DC Discontinue
                  ED Edit
                                                  AL Activity Logs
HD Hold
                       RN Renew
FL Flag VF Verify Select Item(s): Next Screen// AL Activity Logs
  1 - Short Activity Log
  2 - Long Activity Log
  3 - Dispense Log
  4 - History Log
Select LOG to display: 2 Long Activity Log
   Date: 09/07/00 14:07 User: PSJPHARMACIST, ONE
Activity: ORDER VERIFIED BY PHARMACIST
    Date: 09/07/00 14:07
                            User: PSJPHARMACIST, ONE
Activity: ORDER VERIFIED
  Field: Requested Start Date
Old Data: 09/07/00 09:00
   Date: 09/07/00 14:07
                           User: PSJPHARMACIST, ONE
Activity: ORDER VERIFIED
  Field: Requested Stop Date
Old Data: 09/07/00 24:00
Enter RETURN to continue or '^' to exit:
```

4.1.5.7. Finish

When an order is placed or renewed by a provider through CPRS, the nurse or pharmacist needs to accept, finish, and/or verify this order. The same procedures are followed to finish the renewed order as to finish a new order with the following exceptions:

The PENDING RENEWAL orders may be speed finished. The user may enter an **F**, for finish, at the "Select ACTION or ORDERS:" prompt and then select the pending renewals to be finished. A prompt is issued for the Stop Date/Time. This value is used as the Stop Date/Time for the pending renewals selected. All other fields will retain the values from the renewed order.

Note: Order Checks happen during the finish process – refer to the <u>Notes and Screen Example</u> below.

When an action of FN (Finish) is taken on one child order that is part of a Complex Order, a message will display informing the user that the order is part of a Complex Order, and the user is prompted to confirm that the action will be taken on all of the associated child orders.

Note: Complex orders cannot be speed finished because it may not be appropriate to assign the same stop date to all components of a complex order.

Example: Complex Unit Dose Orders with Overlapping Administration Times

When finishing (FN) a complex unit dose drug order with overlapping admin times, after you select the order, a warning message is displayed with the warning and the overlapping admin times.

```
**WARNING**
The highlighted admin times for these portions of this complex order overlap.

Part 1 has a schedule of BID and admin time(s) of 10-22.

AND
Part 2 has a schedule of QDAY and admin time(s) of 10.

Please ensure the schedules and administration times are appropriate.

Press Return to continue...

Enter ?? for more actions

PI Patient Information SO Select Order

PU Patient Record Update NO New Order Entry

Select Action: Next Screen//
```

To finish the order, you must correct the order so that there are no overlapping admin times.

Example: Finish an Order

PENDING UNIT DOSE (ROUTINE) Feb 25, 2001@21:	7:08 Page: 1 of 2
PSJPATIENT1, ONE Ward: 1 EAST	
PID: 000-00-0001 Room-Bed: B-12	Ht(cm):()
DOB: 08/18/20 (80)	Wt(kg):()
*(1)Orderable Item: MULTIVITAMINS TAB	
Instructions: 1 TABLET	
*(2)Dosage Ordered: 1 TABLET	
Duration:	(3)Start: 02/26/01 14:40
*(4) Med Route: PO	REQUESTED START: 02/26/01 14:40
	(5) Stop: 02/28/01 24:00
(6) Schedule Type: FILL on REQUEST	
*(8) Schedule: QDAILY	
(9) Admin Times: 1440	
*(10) Provider: PSJPROVIDER,ONE [es]	
(11) Special Instructions:	
(12) Dispense Drug	U/D Inactive Date
MULTIVITAMIN TABLETS	1
+ Enter ?? for more actions	
BY Bypass FL Flag	
DC Discontinue FN Finish	

Note: When finishing an order, if CPRS Order Checks/Provider Overrides and Pharmacist Interventions exist, they will display upon finishing the order. Heading information displays first, followed by a summary of the Current CPRS Order Checks overridden by the Provider, as well as the Overriding Provider, plus title, Override Entered By, plus title, Date/Time Entered, and the Override Reason.

Example: Finish an Order with Provider Overrides/Interventions

Note: If no Current CPRS Provider Overrides were entered at the time the order was created in CPRS, they will NOT display during finishing, and no heading or messages will display when finishing the Pending order in Inpatient Medications.

The "Action (PBS)" prompt will appear next, with all of the valid actions listed in parentheses. The following are the codes for the possible actions:

- **P** Print specified number of labels now.
- **B** Bypass any more action (entering a caret (^) will also do this).
- S Suspend a specified number of labels for the IV room to print on demand.

The S will only appear as a valid action if the USE SUSPENSE FUNCTIONS site parameter is answered with 1 or YES. The user can perform more than one action, but each action must be done one at a time. As each action is taken, those that operate on labels will reduce the total labels by that amount (e.g., eight labels are needed, three are suspended, then five are available to print).

4.2.3.3. Detailed Allergy/ADR List

The Detailed Allergy/ADR List action displays a detailed listing of the selected item from the patient's Allergy/ADR List. Entry to the *Edit Allergy/ADR Data* option is provided with this list also.

• Enter/Edit Allergy/ADR Data

Provides access to the Adverse Reaction Tracking (ART) package to allow entry and/or edit of allergy adverse reaction data for the patient. See the Allergy package documentation for more information on Allergy/ADR processing.

• **Select Allergy** Allows the user to view a specific allergy.

4.2.3.4. Intervention Menu

This option is only available to those users who hold the PSJ RPHARM key.

The Intervention Menu action allows entry of new interventions and edit, delete, view, or printing of an existing intervention. Each kind of intervention will be discussed and an example will follow.

Note: Interventions can also be dynamically created in response to Order Checks for critical drug-drug interactions and allergy/ADRs. Refer to Section 4.3 Order Checks.

If a change is made to an intervention associated to an inpatient order made in response to Critical Drug-Drug and/or Allergy/ADR, the changes are reflected and displayed whenever interventions display.

New interventions entered via the Intervention Menu are at the patient level and are not associated with a particular order. Consequently, new entries made through this menu are not reflected in the OCI listing, the BCMA Display Order detail report, and do not cause highlighting in BCMA.

New: This option is used to add an entry into the APSP INTERVENTION file.

Example: New Intervention

```
Patient Information Sep 22, 2000 08:03:07 Page: 1 of 1
                                Ward: 1 West
PSJPATIENT2, TWO

        JPATIENT2, TWO
        Ward: 1 West

        PID: 000-00-0002
        Room-Bed: A-6

                                                       Ht(cm): 167.64 (04/21/99)
   DOB: 02/22/42 (58)
                                                      Wt(kg): 85.00 (04/21/99)
   Sex: MALE
                                                    Admitted: 09/16/99
                          Last transferred: ******
   Dx: TEST PATIENT
Allergies - Verified: CARAMEL, CN900, LOMEFLOXACIN, PENTAMIDINE, PENTAZOCINE,
                         CHOCOLATE, NUTS, STRAWBERRIES, DUST
         Non-Verified: AMOXICILLIN, AMPICILLIN, TAPE, FISH,
                         FLUPHENAZINE DECANOATE
               Remote:
   Adverse Reactions:
 Inpatient Narrative: Inpatient narrative
Outpatient Narrative: This is the Outpatient Narrative. This patient doesn't
like waiting at the pickup window. He gets very angry.
           Enter ?? for more actions
PU ratient Record Update NO New Order Entry
DA Detailed Allergy/ADR List IN Intervention Menu
VP View Profile
Select Action: View Profile// IN Intervention Menu
                             --- Intervention Menu ---
    Delete Pharmacy Intervention
Edit Pharmacy Intervention
                                                Print Pharmacy Intervention
DI
                                           PO
                                          VP View Pharmacy Intervention
ED
NE Enter Pharmacy Intervention
Select Item(s): NE
                     Enter Pharmacy Intervention
Select APSP INTERVENTION INTERVENTION DATE: T SEP 22, 2000
  Are you adding 'SEP 22, 2000' as a new APSP INTERVENTION (the 155TH)? No// Y
  (Yes)
   APSP INTERVENTION PATIENT: PSJPATIENT2,TWO 02-22-42 000000002
SC VETERAN
   APSP INTERVENTION DRUG: WAR
     1 WARFARIN 10MG
2 WARFARIN 10MG U/D BL100
3 WARFARIN 2.5MG BL100
4 WARFARIN 2.5MG U/D BL100
BL100TAB
BL100TAB
                                                 TAB
                                    BL100
                                                   TAB **AUTO STOP 2D**
4 WARFARIN 2.5MG U/D BL100 TA
5 WARFARIN 2MG BL100TAB
Press <RETURN> to see more, '^' to exit this list, OR
                                                       TAB **AUTO STOP 2D**
CHOOSE 1-5: 1 WARFARIN 10MG BL100
PROVIDER: PSJPROVIDER, ONE
                                   PROV
INSTITUTED BY: PHARMACY// <Enter> PHARMACY
INTERVENTION: ALLERGY
RECOMMENDATION:
                  NO CHANGE
WAS PROVIDER CONTACTED: N NO
RECOMMENDATION ACCEPTED: Y YES
FINANCIAL COST: <Enter>
REASON FOR INTERVENTION:
 1>
ACTION TAKEN:
 1>
CLINICAL IMPACT:
FINANCIAL IMPACT:
  1>
```

4.2.4.2. Edit

This action allows modification of any field shown on the order view that is preceded by a number in parenthesis (#).

Example: Edit an Order

```
ACTIVE IV
                               Mar 20, 2001@16:41:14 Page: 1 of
PSJPATIENT1, ONE
PID: 000-00-0001
                              Ward: 1 EAST
Room-Bed: B-12
                                                     Ht.(cm):
   DOB: 08/18/20 (80)
                                                     Wt(kg):
*(1) Additives:
                              Order number: 64
                                                          Type: PIGGYBACK
       AMPICILLIN 1000 MG
 (2) Solutions:
       0.9% SODIUM CHLORIDE 100 MT.
                                                * (4)
                                                         Start: 03/19/01 11:30
           Duration:
 (3) Infusion Rate: INFUSE OVER 30 MIN.
                                                * (6)
                                                          Stop: 03/20/01 24:00
*(5) Med Route: IVPB
      Schedule: QID
Admin Times: 09-13-17-21
                                                 Last Fill: 03/19/01 14:57
* (7)
 (8)
                                                     Quantity: 2
          Provider: PSJPROVIDER,ONE [es]
                                                   Cum. Doses: 9
*(10)Orderable Item: AMPICILLIN INJ
      Instructions:
 (11) Other Print:
          Enter ?? for more actions
DC Discontinue ED Edit
HD Hold RN Renew
FL Flag OC On Ca
                                                      AL Activity Logs
                                Renew
                         OC On Call
FL Flag
Select Item(s): Next Screen// ED Edit
Select FIELDS TO EDIT: 11
OTHER PRINT INFO: TESTING
```

If a field marked with an asterisk (*) to the left of the number is changed, the original order will be discontinued, and a new order containing the edited data will be created. The Stop Date/Time of the original order will be changed to the date/time the new edit order is accepted. The old and new orders are linked and may be viewed using the History Log function. When the screen is refreshed, the message, "This change will cause a new order to be created," will be displayed.

Editing Orders with CPRS Overrides/Pharmacist Interventions

The first time a field marked with an asterisk (*), is selected for editing, if CPRS Provider Overrides and/or Pharmacy Interventions exist for the order, entering Y (Yes) at the prompt: "Order Check Overrides/Interventions exist for this order. Display? (Y/N)? Y//" displays the following:

Heading information, followed by a summary of the Current CPRS Order Checks overridden by the Provider, as well as the Overriding Provider, and title, Override Entered By, and title, Date/Time Entered, and the Override Reason.

Refer to "Edit an Order with Provider Overrides/Interventions" for an example of the screen.

Once a Complex Order is made active, the following fields may not be edited:

- ADMINISTRATION TIME
- Any field where an edit would cause a new order to be created. These fields are denoted with an asterisk in the Detailed View of a Complex Order.

If a change to one of these fields is necessary, the Complex Order must be discontinued and a new Complex Order must be created.

Example: Edit an Order (continued)

```
Mar 20, 2001@16:42:02 Page: 1 of 2
ACTIVE IV
PSJPATIENT1, ONE
                           Ward: 1 EAST
  PID: 000-00-0001
                         Room-Bed: B-12
                                            Ht(cm):
  DOB: 08/18/20 (80)
                                            Wt(kg):___
*(1) Additives:
                        Order number: 64
                                               Type: PIGGYBACK
     AMPICILLIN 1000 MG
 (2) Solutions:
     0.9% SODIUM CHLORIDE 100 ML
                                        *(4) Start: 03/19/01 11:30
         Duration:
 (3) Infusion Rate: INFUSE OVER 0 MIN.
     Med Route: IVPB
* (5)
                                        *(6) Stop: 03/20/01 24:00
* (7)
        Schedule: QID
                                          Last Fill: 03/19/01 14:57
    Admin Times: 09-13-17-21
(8)
                                            Quantity: 2
* (9)
      Provider: PSJPROVIDER,ONE [es]
                                          Cum. Doses: 9
*(10)Orderable Item: AMPICILLIN INJ
     Instructions:
 (11) Other Print: TESTING
(12) Remarks :
       Enter ?? for more actions
AC Accept ED Edit
Select Item(s): Next Screen// AC Accept
Orderable Item: MULTIVITAMINS INJ
Give: IVPB QID
[64]0001 1 EAST 03/20/01
PSJPATIENT1, ONE B-12AMPICILLIN 1000 MG 0.9% SODIUM CHLORIDE 100 ML
INFUSE OVER 30 MIN.
TESTING
QID
09-13-17-21
1[1]
Start date: MAR 19,2001 11:30 Stop date: MAR 20,2001 24:00
Is this O.K.: Y// <Enter>
REASON FOR ACTIVITY: <Enter>
7 Labels needed for doses due at \dots
03/19/01 1300 : 03/19/01 1700 : 03/19/01 2100 : 03/20/01 0900 : 03/20/01 1300 :
03/20/01 1700 : 03/20/01 2100 :
                     9
                            12
                                    15
                                            18
                                                   21
            6
^
                                          Ν
Next delivery time is 1500 ***
Action (PB) P// BYPASS
```

4.2.4.5. Renew

Medication orders (referred to in this section as orders) that may be renewed include the following:

- All non-complex active Unit Dose and IV orders.
- Orders that have been discontinued due to ward transfer or treating specialty change.
- Expired orders containing an administration schedule (Unit Dose and scheduled IV orders) that have not had a scheduled administration time since the last BCMA action was taken.
- Expired orders not containing an administration schedule (continuous IV orders) that have had an expired status less than the time limit defined in the EXPIRED IV TIME LIMIT field in the PHARMACY SYSTEM file.

Note: Complex Orders may only be renewed if all associated child orders are renewable.

Renewing Orders with CPRS Overrides/Pharmacist Interventions

When renewing an order, if CPRS Provider Overrides and/or Pharmacy Interventions exist for the order, entering Y (Yes) at the prompt: "Order Check Overrides/Interventions exist for this order. Display? (Y/N)? Y//" displays the following:

Heading information first, followed by a summary of the Current CPRS Order Checks overridden by the Provider, as well as the Overriding Provider, and title, Override Entered By, and title, Date/Time Entered, and the Override Reason.

Refer to "Renew an Order with Provider Overrides/Interventions" for an example of the screen.

Renewing Active Orders

The following applies when the RN (Renew) action is taken on any order with a status of "Active":

- A new Default Stop Date/Time is calculated for the order using the same calculation applied to new orders. The starting point of the Default Stop Date/Time calculation is the date and time that the order was signed in CPRS or the date and time that the RN (Renew) action was taken in Inpatient Medications.
- The RN (Renew) action does not create a new order.
- The Start Date/Time is not available for editing when an order is renewed.

Note: Orders having a schedule type of One-Time or On Call must have a status of "Active" in order to be renewed.

Renewing Discontinued Orders

IV and Unit Dose orders that have been discontinued, either through the (DC) Discontinue action or discontinued due to edit, cannot be renewed.

IV and Unit Dose medication orders that have been discontinued due to ward transfer or treating specialty change will allow the (RN) Renew action.

Renewing Expired Unit Dose Orders

The following applies to expired Unit Dose orders having a schedule type of Continuous or PRN.

- 1. The RN (Renew) action will not be available on an order with a status of "Expired" if either of the following two conditions exist:
 - a. If the difference between the current system date and time and the last scheduled administration time is greater than the frequency of the schedule. This logic will be used for schedules with standard intervals (for example, Q7H).
 - b. If the current system date and time is greater than the time that the next dose is due. This logic is used for schedules with non-standard intervals (for example, Q6H 0600-1200-1800).
- 2. A new Default Stop Date/Time is calculated for the order using the same calculation applied to new orders. The starting point of the Default Stop Date/Time calculation is the date and time that the order was signed in CPRS or the date and time that the RN (Renew) action was taken in Inpatient Medications.
- 3. The (RN) Renew action does not create a new order.
- 4. The Start Date/Time is not available for editing when an order is renewed.
- 5. The renewed order has a status of "Active."

Example: Activity Log (continued)

```
*(3) Infusion Rate: 80 ml/hr
                                         * (6)
     Med Route: IV
                                                  Stop: 02/20/02 24:00
* (5)
BCMA ORDER LAST ACTION: 02/20/02 15:50 Infusing*
*(7) Schedule:
                                              Last Fill: 02/20/02 15:55
(8) Admin Times:
*(9) Provider: PSJPROVIDER,ONE [es]
(10) Other Print:
                                               Quantity: 1
                                              Cum. Doses: 1
(11) Remarks :
        Enter ?? for more actions
DC Discontinue RN Renew
HD Hold OC On Call
ED Edit AL Activity Logs
Select Item(s): Next Screen// AL Activity Logs
(A) ctivity (L) abel (H) istory: Activity Log
ACTIVITY LOG:
# DATE TIME REASON
1 FEB 20,2002 15:55:09 COMPLETE
                                             PSJPHARMACIST, ONE
  Comment: DISCONTINUED (EDIT)
2 FEB 20,2002 15:55:12 VERIFY
                                              PSJPHARMACIST, ONE
Comment: ORDER VERIFIED BY PHARMACIST
(A) ctivity (L) abel (H) istory: Label Log
LABEL LOG:
# DATE/TIME ACTION USER
                                   #LABELS TRACK
_______
1 FEB 20,2002@15:55:12
                DISPENSED PSJPHARMACIST, ONE 1 ORDER ACTION
Enter RETURN to continue or '^' to exit: <Enter>
Unique IDs for this order:
Label Date/Time Unique ID Status Count BCMA Action - Date/Time
02/20/02 15:55 739V443
(A) ctivity (L) abel (H) istory: History Log
DEVICE: HOME// <Enter> NT/Cache virtual TELNET terminal Right Margin: 80// <Enter>
    Patient: PSJPATIENT4, FOUR
                                                  Status: DISCONTINUED
*(1) Additives: Order number: 444
                                                  Type: ADMIXTURE
      POTASSIUM CHLORIDE XXXXXXXXXX 35 MEQ
*(2) Solutions:
     DEXTROSE 5% 1/2 NS 1000 ML
                                         *(4) Start: 02/20/02 15:46
*(4) Start: 02/20/02 15:46
*(6) Stop: 02/20/02 15:55
         Duration:
*(3) Infusion Rate: 80 ml/hr
* (5)
      Med Route: IV
BCMA ORDER LAST ACTION: 02/20/02 15:50 Infusing
                                              Last Fill: 02/20/02 15:46
*(7) Schedule: (8) Admin Times:
                                               Quantity: 2
* (9)
      Provider: PSJPROVIDER, ONE [es] Cum. Doses: 2
(10) Other Print:
 (11) Remarks:
     Entry By: PSJPROVIDER, ONE
                                              Entry Date: 02/20/02 15:46
Enter RETURN to continue or '^' to exit: <Enter>
```

January 2005

Example: Activity Log (continued)

```
Patient: PSJPATIENT4, FOUR
                                                    Status: ACTIVE
                   Order number: 445
*(1) Additives:
                                                      Type: ADMIXTURE
      POTASSIUM CHLORIDE XXXXXXXXXX 35 MEQ
*(2) Solutions:
      DEXTROSE 5% 1/2 NS 1000 ML
                                            *(4) Start: 02/20/02 15:46
Duration:
*(3) Infusion Rate: 80 ml/hr
         Duration:
                                           * (6)
                                                     Stop: 02/20/02 24:00
*(5) Med Route: IV
BCMA ORDER LAST ACTION: 02/20/02 15:50 Infusing*
* (7)
        Schedule:
                                                Last Fill: 02/20/02 15:55
(8) Admin Times:
*(9) Provider:
(10) Other Print:
                                                 Quantity: 1
        Provider: PSJPROVIDER,ONE [es]
                                                Cum. Doses: 1
 (11) Remarks:
      Entry By: PSJPROVIDER, ONE
                                              Entry Date: 02/20/02 15:55
Enter RETURN to continue or '^' to exit: <Enter>
(A) ctivity (L) abel (H) istory:
```

4.2.4.7. Finish

Users must hold the PSJ RPHARM key for the ability to finish orders placed through CPRS.

Pharmacy Technicians must hold the PSJI PHARM TECH key for the ability to finish orders placed through CPRS. These users are not allowed to verify orders, only finish orders.

When an order is placed or renewed by a provider through CPRS, the pharmacist needs to finish this order. The same procedures are followed to finish the renewed order as to finish a new order.

When an action of FN (Finish) is taken on one child order that is part of a Complex Order, a message will display informing the user that the order is part of a Complex Order, and the user is prompted to confirm that the action will be taken on all of the associated child orders.

Note: Complex orders cannot be speed finished because it may not be appropriate to assign the same stop date to all components of a complex order.

Note: When finishing an order, if CPRS Order Checks/Provider Overrides and Pharmacist Interventions exist, they will display during the finish process. Heading information displays first, followed by a summary of the Current CPRS Order Checks overridden by the Provider, as well as the Overriding Provider, and title, Override Entered By, and title, Date/Time Entered, and the Override Reason. Refer to "*Finish an Order with Provider Overrides/Interventions*" for an example of the screen.

Note: For more details on ordering, see *New Order Entry*.

4.2.4.8. Expected First Dose Changes

Inpatient Medications no longer displays an expected first dose for an order containing a schedule with a schedule type of One-time. The system also no longer displays an expected first dose for an order containing a schedule with a schedule type of On-call.

(This page included for two-sided copying.)

```
Display Interaction Monograph? No//
***Significant*** Drug Interaction with Prospective Drug:
                   SIMVASTATIN 10MG TAB and
        Local RX#: 504361
              Drug: WARFARIN (COUMADIN) NA 5MG TAB (Active)
              SIG: TAKE ONE TABLET BY MOUTH EVERY 2 HOURS
     Processing Status: Not released locally (Window)
     Pending Order: WARFARIN (COUMADIN) NA 5MG TAB
              SIG: TAKE ONE TABLET BY MOUTH EVERY 2 HOURS
  *** Refer to MONOGRAPH for SIGNIFICANT INTERACTION CLINICAL EFFECTS
Display Interaction Monograph? No//
***Significant*** Drug Interaction with Prospective Drug:
                   SIMVASTATIN 10MG TAB and
     Pending Order: RIFAMPIN 300MG CAP
              SIG: TAKE ONE CAPSULE BY MOUTH EVERY DAY
  *** Refer to MONOGRAPH for SIGNIFICANT INTERACTION CLINICAL EFFECTS
Display Interaction Monograph? No// NO
Do you want to Continue? Y// NO
RX DELETED
```

• **Drug-Drug Interactions** - Drug-drug interactions will be either critical or significant. If the Dispense Drug selected is identified as having an interaction with one of the drugs the patient is already receiving, the order the new drug interacts with will be displayed.

Note: For a Significant Interaction, the user who holds the PSJ RPHARM key is allowed to enter an intervention, but one is not required. For a Critical Interaction, the user who holds the PSJ RPHARM key must enter an intervention before continuing.

Note: If the user (who holds the PSJ RPHARM key), is prompted for an intervention and enters 9, which is OTHER, "OTHER FOR RECOMMENDATION" displays. This allows the user to enter unlimited free text as a response to the order check(s).

Example: Drug-Drug Interaction Display

```
Mar 17, 2011@10:40
Patient Information
                                                             Page:
                                                                       1 of
                             Ward: 7A GEN A
Room-Bed: Ht(cm): 175.26 (12/15/08)
Wt(kg): 100.00 (12/15/08)
Admitted: 01/31/02
BCMA, EIGHTEEN-PATIENT
   PID: 666-33-0018
DOB: 04/07/35 (75)
   Sex: FEMALE
                                                   Admitted: 01/31/02
   Dx: UPSET
                                         Last transferred: 06/04/10
Allergies - Verified: AMPICILLIN, PENICILLIN, STRAWBERRIES
        Non-Verified:
   Adverse Reactions:
 Inpatient Narrative:
Outpatient Narrative:
          Enter ?? for more actions
PU Patient Record Update
                            NO New Order Entry
```

```
DA Detailed Allergy/ADR List IN Intervention Menu
VP View Profile
Select Action: View Profile// View Profile
SHORT, LONG, or NO Profile? SHORT// SHORT
Inpatient Order Entry Mar 17, 2011@10:40:12 Page: 1 of 2

BCMA, EIGHTEEN-PATIENT Ward: 7A GEN

PID: 666-33-0018 Room-Bed: Ht(cm): 175.26 (12/15/08)

DOR: 04/07/35 (75) Wt (kg): 100.00 (12/15/08)
                                                         Wt(kg): 100.00 (12/15/08)
   DOB: 04/07/35 (75)
                                                  Admitted: 01/31/02
   Sex: FEMALE
   Dx: UPSET
                                          Last transferred: 06/04/10
                                                  C 03/16 03/17 A
  1 INDINAVIR CAP, ORAL
         Give: 400MG PO QDAY
   2 SIMVASTATIN TAB
                                                   C 03/16 03/18 A
         Give: 40MG PO QPM
         ----NON-VERIFIEDCOMPLEX------
   3 LITHIUM TAB, SA
                                                   C 10/13 10/15 N
          Give: 450MG PO QID
        LITHIUM TAB, SA
                                                  C 10/13 10/15 N
         Give: 10000MG PO Q4H
      RILUZOLE TAB
                                                  C 10/13 10/15 N
        Give: 50MG PO BID
          Enter ?? for more actions
PI Patient Information SO Select Order
PU Patient Record Update NO New Order En
PU Patient Record Update
                                         NO New Order Entry
Select Action: Next Screen// NO New Order
Eligibility: SERVICE CONNECTED 50% to 100% SC%: 60 RX PATIENT STATUS: SC//
DRUG: WARFAR
  Lookup: GENERIC NAME
    1 WARFARIN (COUMADIN) 5MG INJ
2 WARFARIN (COUMADIN) NA 1MC MAR
                                                BL110
         WARFARIN (COUMADIN) NA 1MG TAB
     2 WARFARIN (COUMADIN) NA 1MG TAB BL110
3 WARFARIN (COUMADIN) NA 5MG TAB BL110
     4 WARFARIN (COUMADIN) NA 7.5MG TAB BREKSEN
5 WARFARIN (COUMADIN) NA 10MG TAB BL110
Press <RETURN> to see more, '^' to exit this list, '^^' to exit all lists, OR
CHOOSE 1-5: 1 WARFARIN (COUMADIN) 5MG INJ
Now doing allergy checks. Please wait...
Now Processing Enhanced Order Checks! Please wait...
Order Checks could not be done for Drug: BACLOFEN 10MG TAB, please
complete a manual check for Drug Interactions, Duplicate Therapy and
appropriate Dosing.
Press Return to Continue:
***Critical*** Drug Interaction with Prospective Drug:
                     WARFARIN (COUMADIN) 5MG INJ and
         Local RX#: 504196
               Drug: AMIODARONE HCL (PACERONE) 200MG TAB (Active)
                SIG: TAKE THREE TABLETS BY MOUTH EVERY 3 HOURS
     Processing Status: Not released locally (Mail)
     Pending Order: AMIODARONE HCL (PACERONE) 200MG TAB
               SIG: TAKE THREE TABLETS BY MOUTH EVERY 3 HOURS
        Non-VA Med: AMIODARONE HCL (PACERONE) 200MG TAB
            Dosage: 400MG Schedule: EVERY DAY
```

The concurrent administration of amiodarone and an anticoagulant may result in an increase in the clinical effects of the anticoagulant and an increased risk of bleeding.(1-22) It may take several weeks of concurrent therapy before the full effects of this interaction are noted. The effect of amiodarone on anticoagulant levels may continue for several months after amiodarone is discontinued.

Display Interaction Monograph? No// NO

Press return to continue:

Critical Drug Interaction with Prospective Drug: WARFARIN (COUMADIN) 5MG INJ and

Local RX#: 504183

Drug: CIMETIDINE 300MG TAB (Active)

SIG: TAKE TWO TABLETS BY MOUTH EVERY 3 HOURS

Processing Status: Not released locally (Mail)

Pending Order: CIMETIDINE 300MG TAB

SIG: TAKE TWO TABLETS BY MOUTH EVERY 3 HOURS

The pharmacologic effects of warfarin may be increased resulting in severe

bleeding.

Display Interaction Monograph? No// NC

Critical Drug Interaction with Prospective Drug: WARFARIN (COUMADIN) 5MG INJ and

Pending Order: RIFAMPIN 300MG CAP

SIG: TAKE ONE CAPSULE BY MOUTH EVERY DAY

Concurrent or recent use of a rifamycin may result in decreased levels of and clinical effects from anticoagulants. If the rifamycin is withdrawn, levels and effects of the anticoagulant may increase, increasing the risk of hemorrhage. This effect may be dose-related and continue beyond discontinuation of the rifamycin.

Display Interaction Monograph? No// NO

Significant Drug Interaction with Prospective Drug: WARFARIN (COUMADIN) 5MG INJ and

Local RX#: 504280

Drug: INDINAVIR SULFATE 400MG CAP (Active) SIG: TAKE ONE CAPSULE BY MOUTH EVERY DAY Processing Status: Not released locally (Window)

*** Refer to MONOGRAPH for SIGNIFICANT INTERACTION CLINICAL EFFECTS

Display Interaction Monograph? No// NO

Significant Drug Interaction with Prospective Drug: WARFARIN (COUMADIN) 5MG INJ and

Local RX#: 504426

Drug: SIMVASTATIN 40MG TAB (Suspended)

SIG: TAKE 20 TABLETS BY MOUTH EVERY 4 HOURS AND TAKE 15
TABLETS TWICE A DAY BEFORE MEALS AND TAKE TEN TABLETS
TWICE A DAY AND TAKE FIVE TABLETS EVERY 3 HOURS AND
TAKE ONE SIXTY MG TABLET(S) Q5H AND TAKE ONE FORTY MG

TABLET(S) EVERY EVENING PAT INSTRUCTIONS

```
Processing Status: Not released locally (Mail)

*** Refer to MONOGRAPH for SIGNIFICANT INTERACTION CLINICAL EFFECTS

Display Professional Interaction Monograph(s)? NO//

Do you want to Continue with METRONIDAZOLE 500MG IN 100ML? NO// YES

Now creating Pharmacy Intervention
For METRONIDAZOLE 500MG IN 100ML

PROVIDER: PROV INPATIENT-MEDS, PROVIDER PROV
RECOMMENDATION: 9 OTHER
OTHER FOR RECOMMENDATION:
No existing text
Edit? NO// YES

==[ WRAP ]==[ INSERT ]======< OTHER FOR RECOMMENDATION >=====[ <PF1>H=Help ]====
Discussed with doctor, ok to administer.
```

Note: The "OTHER FOR RECOMMENDATION" text field is best used for the Pharmacist reason for overriding the order check(s). For critical drug-drug and allergy/ADR interactions, this information will display when the OCI 'Hidden Action' is used in Inpatient Medications. It will also be available for the nurse to view in the BCMA Display Order detail report.

• **Drug-Allergy Interactions** – If the Dispense Drug selected is identified as having an interaction with one of the patient's allergies, the allergy the drug interacts with will be displayed. Pharmacist Interventions for Drug-Allergy/ADR Interactions are optional.

Note: If the user (who holds the PSJ RPHARM key), is prompted for an intervention and enters 9, which is OTHER, "OTHER FOR RECOMMENDATION" displays. This allows the user to enter unlimited free text as a response to the order check(s).

Example: Remote Allergy/ADR - New Order Entry Backdoor - Both Ingredient and Drug Class Defined

```
Select Action: View Profile// NO New Order Entry
Select DRUG: DILTIAZEM
  Lookup: GENERIC NAME
    1 DILTIAZEM (INWOOD) 120MG SA CAP
                                                    CV200
     2 DILTIAZEM (INWOOD) 180MG SA CAP
3 DILTIAZEM (INWOOD) 240MG SA CAP
                                                         CV200
    3 DILTIAZEM (INWOOD) 240MG SA CAP
4 DILTIAZEM (INWOOD) 300MG SA CAP CV200
CV200
                                                         CV200
    5 DILTIAZEM (INWOOD) 360MG SA CAP
Press <RETURN> to see more, '^' to exit this list, '^^' to exit all lists, OR CHOOSE 1-5: 1 DILTIAZEM (INWOOD) 120MG SA CAP CV200
CHOOSE 1-5: 1 DILTIAZEM (INWOOD) 120MG SA CAP
A Drug-Allergy Reaction exists for this medication and/or class!
Drug: DILTIAZEM (DILACOR XR) 240MG SA CAP
   Ingredients: DILTIAZEM (REMOTE SITE(S)),
   Drug Class: CV200 CALCIUM CHANNEL BLOCKERS (REMOTE SITE(S))
```

```
Do you want to Intervene NO// YES

Now creating Pharmacy Intervention
For DILTIAZEM (INWOOD) 120MG SA CAP

PROVIDER: PSJPROVIDER, ONE OP PROVIDER
RECOMMENDATION: 9 OTHER
OTHER FOR RECOMMENDATION:
No existing text
Edit? NO// YES

==[ WRAP ]==[ INSERT ]=====< OTHER FOR RECOMMENDATION >=====[ <PF1>H=Help ]====
Discussed with doctor and okay to administer.
```

Note: The "OTHER FOR RECOMMENDATION" text field is best used for the Pharmacist reason for overriding the order check(s). For critical drug-drug and allergy/ADR interactions, this information will display when the OCI 'Hidden Action' is used in Inpatient Medications. It will also be available for the nurse to view in the BCMA Display Order detail report.

• CPRS Order Check: Aminoglycoside Ordered

```
Aminoglycoside Ordered
Trigger: Ordering session completion.
Mechanism: For each medication order placed during this ordering session, the CPRS DANGEROUS MEDS FOR PT > 64 - Yes
This is based on the BEERS list. This order check only checks for three drugs:
Amitriptyline, Chlorpropamide and Dipyridamole. The workgroup felt that the list of
drugs should be expanded. A request can be sent to CPRS for this.
[Note: The creatinine clearance value displayed in some order check messages is an
estimate based on adjusted body weight if patient height is > 60 inches. Approved by
the CPRS Clinical Workgroup 8/11/04, it is based on a modified Cockcroft-Gault
formula and was installed with patch OR*3*221.
For more information: http://www.ascp.com/public/pubs/tcp/1999/jan/cockcroft.shtml
CrCl (male) = (140 - age) x (adj body weight* in kg)
(serum creatinine) x 72
       * If patient height is not greater than 60 inches, actual body weight is used.
CrCl (female) = 0.85 \times CrCl (male)
To calculate adjusted body weight, the following equations are used:
Ideal body weight (IBW) = 50 \text{ kg x (for men)} or 45 \text{ kg x (for women)} + 2.3 \text{ x (height in }
inches - 60)
Adjusted body weight (Adj. BW) if the ratio of actual BW/IBW > 1.3 = (0.3 x (Actual
BW - IBW)) + IBW
Adjusted body weight if the ratio of actual BW/IBW is not > 1.3 = IBW or Actual BW
(whichever is less)]
Message: Aminoglycoside - est. CrCl: <value calculated from most recent serum
creatinine >. (CREAT: <result > BUN: <result >).
Danger Lvl: This order check is exported with a High clinical danger level.
```

CPRS Order Check: Dangerous Meds for Patients >64

January 2012 Inpatient Medications V. 5.0 124g

Trigger: Acceptance of pharmacy orderable items amitriptyline, chlorpropamide or dipvridamole. Mechanism: The CPRS Expert System determines if the patient is greater than 64 years old. It then checks the orderable item of the medication ordered to determine if it is mapped as a local term to the national term DANGEROUS MEDS FOR PTS > 64. Message: If the orderable item text contains AMITRIPTYLINE this message is displayed: Patient is <age>. Amitriptyline can cause cognitive impairment and loss of balance in older patients. Consider other antidepressant medications on formulary. If the orderable item text contains CHLORPROPAMIDE this message is displayed: Patient is <age>. Older patients may experience hypoglycemia with Chlorpropamide due do its long duration and variable renal secretion. They may also be at increased risk for Chlorpropamide-induced SIADH. If the orderable item text contains DIPYRIDAMOLE this message is displayed: Patient is <age>. Older patients can experience adverse reactions at high doses of Dipyridamole (e.g., headache, dizziness, syncope, GI intolerance.) There is also questionable efficacy at lower doses. Danger Lvl: This order check is exported with a High clinical danger level.

CPRS Order Check: Glucophage Lab Results

Glucophage-Lab Results Interactions
Trigger: Selection of a Pharmacy orderable item.
Mechanism: The CPRS Expert System checks the pharmacy orderable item's local text
(from the Dispense Drug file [#50]) to determine if it contains "Glucophage" or
"metformin". The expert system next searches for a serum creatinine result within the
past x number of days as determined by parameter ORK GLUCOPHAGE CREATININE. If the
patient's creatinine result was greater than 1.5 or does not exist, OE/RR is notified
and the warning message is displayed.

Message: Metformin- no serum creatinine within past <x> days. else:
 Metformin - Creatinine results: <creatinine greater than 1.5 w/in past <x> days>
Danger Lvl: This order check is exported with a High clinical danger level.

1.1.1. Inpatient Duplicate Therapy

Inpatient orders are checked for therapeutic duplication with drugs within the same class. If orders have the same drug (meaning the same class), they will be included in the list. The header for Inpatient Duplicate Therapy will be like: This patient is already receiving the following:

INPATIENT and/or OUTPATIENT order(s) for a drug in the same therapeutic class(es) as SIMVASTATIN 40MG TAB:

The user will have the opportunity to discontinue duplicate order(s) after the banner...

Example: Duplicate Therapy Banner

```
This patient is already receiving the following INPATIENT and/or OUTPATIENT order(s) for a drug in the same therapeutic class(es) as SIMVASTATIN 40MG TAB:

GEMFIBROZIL TAB,ORAL C 02/08 05/19 A

Give: 600MG PO BID

GEMFIBROZIL TAB,ORAL C 02/08 05/19 A

Give: 600MG PO BID

Local Rx #504563 (ACTIVE) for FLUVASTATIN NA 20MG CAP

SIG: TAKE ONE CAPSULE BY MOUTH TWICE A DAY

Processing Status: Not released locally (Window)
```

4.3.3. Allergy/ADR Example Order Checks

Inpatient Medications (Unit Dose and IV) order entry process with check for adverse allergy/ADR reactions: (conditions by which the user will get new order checks)

- Entering a new IV or Unit Dose medication order through pharmacy options
- Finishing a pending IV or Unit Dose medication order
- Renewing an IV or Unit Dose order
- Creating a new Unit Dose order when editing the orderable item (to a new orderable item) through pharmacy options
- When editing the IV additive field (changing existing additive or adding new additive) for an IV order through pharmacy options
- When editing the IV solution field (changing existing solution or adding a new solution) for an IV order through pharmacy options – This applies only to IV solutions marked as a PreMix
- Entering a new Unit Dose medication order through pharmacy options
- using order sets
- Copying an IV or Unit Dose medication order, thereby creating a new order.

Pharmacist Interventions for Allergy/ADR interactions are optional. Only one warning will be displayed for an Allergy/ADR. The Allergy/ADR warning shall display the following information:

- Drug Text 'A Drug-Allergy Reaction exists for this medication and/or class:'
- Drug Name
- Ingredient(s) (Indicate Local and/or Remote sites) if available
- VA Drug Class(es) (Indicate Local and/or Remote sites) if available

More than one ingredient and more than one VA Drug Class may be associated with an Allergy/ADR. After the Allergy/ADR warning is displayed, the system shall prompt the user if they want to intervene. The default for this prompt shall be 'No.' If the user chooses to intervene, the system will proceed with the intervention dialog. If the user chooses not to intervene, the system will proceed with the order entry dialog.

Example: Remote Allergy/ADR - New Order Entry Backdoor - Both Ingredient and Drug Class Defined

```
Select DRUG: DILTIAZEM
Lookup: GENERIC NAME

1 DILTIAZEM (INWOOD) 120MG SA CAP CV200
2 DILTIAZEM (INWOOD) 240MG SA CAP CV200
3 DILTIAZEM (INWOOD) 240MG SA CAP CV200
4 DILTIAZEM (INWOOD) 360MG SA CAP CV200
5 DILTIAZEM (INWOOD) 360MG SA CAP CV200
Press <RETURN> to see more, '^' to exit this list, '^^' to exit all lists, OR CHOOSE 1-5: 1 DILTIAZEM (INWOOD) 120MG SA CAP CV200

A Drug-Allergy Reaction exists for this medication and/or class!

Drug: DILTIAZEM (DILACOR XR) 240MG SA CAP Ingredients: DILTIAZEM (REMOTE SITE(S)),
Drug Class: CV200 CALCIUM CHANNEL BLOCKERS (REMOTE SITE(S))
```

```
Do you want to Intervene NO// YES

Now creating Pharmacy Intervention
For DILTIAZEM (INWOOD) 120MG SA CAP

PROVIDER: PSJPROVIDER, ONE OP PROVIDER
RECOMMENDATION: 9 OTHER
OTHER FOR RECOMMENDATION:
No existing text
Edit? NO// YES

==[ WRAP ]==[ INSERT ]=====< OTHER FOR RECOMMENDATION >=====[ <PF1>H=Help ]====
Discussed with doctor and okay to administer.
```

Note: The "OTHER FOR RECOMMENDATION" text field is best used for the Pharmacist reason for overriding the order check(s). For critical drug-drug and allergy/ADR interactions, this information will display when the OCI 'Hidden Action' is used in Inpatient Medications. It will also be available for the nurse to view in the BCMA Display Order detail report.

```
Example: New Order Entry - Backdoor - Local & Remote Allergy/ADR - Ingredients & Drug Class exist
Select Unit Dose Medications Option: IOE Inpatient Order Entry
You are signed on under the GLRISC IV ROOM
Current IV LABEL device is: TELNET
Current IV REPORT device is: NULL DEVICE
Select PATIENT: PSJPATIENT, TEN
                                          000-00-0000 02/02/39
                                                                     3AS
               *** Patient Requires a Means Test **
VP View Profile
Allergies - Verified: PENICILLIN, ASPIRIN
       Non-Verified: CODEINE PHOSPHATE 15MG TAB, DIAZEPAM, TETRACYCLINE
Reactions - Verified: SULFAMETHOXAZOLE/TRIMETHOPRIM, VANCOMYCIN
       Non-Verified:
Inpatient Narrative: Place All Meds in NS
Outpatient Narrative:
          Enter ?? for more actions
DA Detailed Allergy/ADR List
PU Patient Record Update
                                       NO New Order Entry
                                      IN Intervention Menu
VP View Profile
Select Action: View Profile// NO New Order Entry
Select DRUG: SULFAMET
 Lookup: GENERIC NAME
SULFAMETHOXAZOLE/TRIMETHOPRIM DS TAB
                                             AM650
        ...OK? Yes//
                      (Yes)
A Drug-Allergy Reaction exists for this medication and/or class!
Drug: SULFAMETHOXAZOLE/TRIMETHOPRIM DS TAB
   Ingredients: SULFAMETHOXAZOLE (LOCAL), TRIMETHOPRIM (LOCAL)
   Drug Class: AM650 SULFONAMIDE/RELATED ANTIMICROBIALS (LOCAL AND REMOTE SITE(S))
```

Example: IV New Order Entry Backdoor

```
Select Action: Next Screen// NO New Order Entry
Select IV TYPE: PIGGYBACK.
Select ADDITIVE: CEFAZOLIN
              *N/F*
 Restriction/Guideline(s) exist. Display? : (N/D): No// NO
(The units of strength for this additive are in GM)
Strength: 1 1 GM
Select ADDITIVE:
Select SOLUTION: D5250 5% DEXTROSE 250 ML
 Restriction/Guideline(s) exist. Display? : (N/D/O/B): No// NO
Now Processing Enhanced Order Checks! Please wait...
This patient is already receiving the following INPATIENT and/or OUTPATIENT order(s)
for a drug in the same therapeutic class as CEFAZOLIN 1GM:
CEFOXITIN INJ
                                    C 03/20 04/03 A
Give: 1GM/1VIAL IM Q12H
PENICILLIN TAB
                                    C 03/20 03/27 A
Give: 500MG PO QID
Duplicate Therapy Class(es): Beta-Lactams
______
CEFOXITIN INJ
                                   C 03/20 04/03 A
Give: 1GM/1VIAL IM Q12H
Duplicate Therapy Class(es): Cephalosporins
Do you wish to DISCONTINUE any of the listed INPATIENT orders? NO// Yes
                                      C 03/20 04/03 A
1. CEFOXITIN INJ
  Give: 1GM/1VIAL IM Q12H
2. PENICILLIN TAB
                                     C 03/20 03/27 A
      Give: 500MG PO QID
Select (1-2): 1
CEFOXITIN INJ
                                 C 03/20 04/03 A
Give: 1GM/1VIAL IM Q12H
```

```
Do you want to discontinue this order? Yes// (Yes)

NATURE OF ORDER: WRITTEN// W
Requesting PROVIDER: PROVIDER, ONE// LBB 119

INFUSION RATE: OVER 30 MINTUES

MED ROUTE: IV//IVPB IV PIGGYBACK IVPB
SCHEDULE: Q12H
ADMINISTRATION TIMES: 09-21//
```

1.1.2. Display of Provider Overrides and Pharmacist Interventions

In Inpatient Medications, the first time a field preceded by an asterisk (*) is selected for editing and when renewing an order, if Current Pharmacist Interventions exist for the order, entering Y (Yes) at the prompt, "Order Check Overrides/Interventions exist for this order. Display? (Y/N)? Y//," will display the following information when the fields are populated with data:

- Heading: **Current Pharmacist Interventions for this order**
- Intervention Date/Time
- Provider
- Pharmacist
- Drug,
- Instituted By
- Intervention
- Other For Recommendation
- Originating Package
- Was Provider Contacted
- Provider Contacted
- Recommendation Accepted
- Agree With Provider
- Rx #
- Division
- Financial Cost
- Other For Intervention
- Reason For Intervention
- Action Taken
- Clinical Impact
- Financial Impact

```
ST) 0.5MG(1/2X1MG) TAB [UNRELEASED] - Concurrent use of anticoagulants with
 metronidazole or tinidazole may result in reduced prothrombin activity and/or
 increased risk of bleeding. - Monograph Available
Press RETURN to Continue or '^' to Exit :
______
      ** Current Pharmacist Interventions for this order **
Intervention Date: 7/12/11 09:14
Provider: PSJPROVIDER, ONE
                                      Pharmacist: PSJPHARMACIST, ONE
Drug: METRONIDAZOLE 250MG TAB Instituted By: PHARMACY
Intervention: CRITICAL DRUG INTERACTION
Recommendation: OTHER
                                     Originating Package: INPATIENT
Other For Recommendation:
 INTERVENTION FOR CRITICAL DRUG-DRUG
Press RETURN to Continue or '^' to Exit:
```

Intervention TIME displays to the right of the date (e.g., 01/18/11 09:04)

If Historical Overrides/Interventions exist for an order, entering Y (Yes) at the prompt: "View Historical Overrides/Interventions for this order (Y/N)? Y//," displays the Historical Pharmacist Intervention information:

```
** Historical Pharmacist Interventions for this order **
Intervention Date: 07/12/11 09:14
Provider: PSJPROVIDER, ONE Pharmacist: PSJPHARMACIST, ONE Drug: METRONIDAZOLE 250MG TAB Instituted By: PHARMACY
Intervention: CRITICAL DRUG INTERACTION
Recommendation: OTHER
                                         Originating Package: INPATIENT
Other For Recommendation:
 Testing 9 OTHER
Press RETURN to Continue or '^' to Exit :
     ** Historical Provider Overrides for this order **
______
Overriding Provider: PSJPROVIDER, ONE (PROVIDER)
Override Entered By: PSJPROVIDER, ONE (PROVIDER)
 Date/Time Entered: 07/12/11 09:13
   Override Reason: Testing 9 OTHER
 CRITICAL drug-drug interaction: METRONIDAZOLE 250MG TAB and WARFARIN NA \,
  (GOLDEN STATE) 2MG TAB [ACTIVE] - Concurrent use of anticoagulants with
 metronidazole or tinidazole may result in reduced prothrombin activity and/or
 increased risk of bleeding. - Monograph Available
 CRITICAL drug-drug interaction: METRONIDAZOLE 250MG TAB and WARFARIN(GOLDEN
 ST) 0.5MG(1/2X1MG) TAB [UNRELEASED] - Concurrent use of anticoagulants with
 metronidazole or tinidazole may result in reduced prothrombin activity and/or
  increased risk of bleeding. - Monograph Available
```

Intervention TIME displays to the right of the date (e.g., 01/18/11 09:04. Current Pharmacist Intervention fields and labels also display, when the fields are populated.

Note: In Inpatient Medications, if no Current Pharmacist Interventions exist when editing a field preceded by an asterisk (*),the following displays:

======	=========	=========	==========	==========	==========
	** CLIRRENT PH	ARMACTST INTER\	VENTIONS FOR TH	TS ORDER **	
	OUTILLIAT 1 11/	THILLIE	VENTIONO TON TH	TO ONDEN	
	=========				==========

NO PHARMACIST INTERVENTIONS TO DISPLAY

2. Glossary

Action Prompts

There are three types of Inpatient Medications "Action" prompts that occur during order entry: ListMan, Patient/Order, and Hidden action prompts.

ListMan Action Prompts

+ Next Screen

- Previous Screen

UP Up a Line
DN Down a Line

> Shift View to Right

< Shift View to Left

FS First screen
LS Last Screen
GO Go to Page

RD Re Display Screen

PS Print Screen
PT Print List
SL Search List

Q Quit

ADPL Auto Display (on/off)

Patient/Order Action Prompts

PU Patient Record Updates

DA Detailed Allergy/ADR List

VP View Profile

NO New Orders Entry
IN Intervention Menu
PI Patient Information

SO Select Order

DC Discontinue

ED Edit
FL Flag
VF Verify
HD Hold

Patient/Order Action Prompts

(continued) RN Renew

AL Activity Logs

OC On Call

NL Print New IV LabelsRL Reprint IV Labels

RC Recycled IV
DT Destroyed IV
CA Cancelled IV

Hidden Action Prompts

LBL Label Patient/Report

JP Jump to a Patient

OTH Other Pharmacy Options

MAR MAR Menu

DC Speed Discontinue

RN Speed Renew SF Speed Finish SV Speed Verify

CO Copy

N Mark Not to be Given

I Mark Incomplete
DIN Drug Restr/Guide

OCI Order Check/Interventions

Active Order Any order which has not expired or been discontinued.

Active orders also include any orders that are on hold or

on call.

Activity Reason Log The complete list of all activity related to a patient

order. The log contains the action taken, the date of the

action, and the user who took the action.

Activity Ruler The activity ruler provides a visual representation of the

relationship between manufacturing times, doses due, and order start times. The intent is to provide the onthe-floor user with a means of tracking activity in the IV room and determining when to call for doses before the normal delivery. The activity ruler can be enabled or disabled under the *SIte Parameters (IV)* option.

CLINIC DEFINITION File

File #53.46. This file is used in conjunction with Inpatient Medications for Outpatients (IMO) to give the user the ability to define, by clinic, default stop dates, whether to auto-dc IMO orders, and whether to send IMO orders to BCMA.

Clinic Group

A clinic group is a combination of outpatient clinics that have been defined as a group within Inpatient Medications to facilitate processing of orders.

Complex Order

An order that is created from CPRS using the Complex Order dialog and consists of one or more associated Inpatient Medication orders, known as "child" orders. Inpatient Medications receives the parent order number from CPRS and links the child orders together. If an action of FN (Finish), VF (Verify), DC (Discontinue), or RN (Renew) is taken on one child order, the action must be taken on all of the associated child orders. For example:

- If one child order within a Complex Order is made active, all child orders in the Complex Order must be made active.
- If one child order within a Complex Order is discontinued, all child orders in the Complex Order must be discontinued.
- If one child order within a Complex Order is renewed, all child orders in the Complex Order must be renewed.

Continuous IV Order

Inpatient Medications IV order not having an administration schedule. This includes the following IV types: Hyperals, Admixtures, Non-Intermittent Syringe, and Non-Intermittent Syringe or Admixture Chemotherapy.

Continuous Syringe

A syringe type of IV that is administered continuously to the patient, similar to a hyperal IV type. This type of syringe is commonly used on outpatients and administered automatically by an infusion pump.

Coverage Times The start and end of coverage period designates

administration times covered by a manufacturing run. There must be a coverage period for all IV types: admixtures and primaries, piggybacks, hyperals, syringes, and chemotherapy. For one type, admixtures for example, the user might define two coverage

periods; one from 1200 to 0259 and another from 0300

to 1159 (this would mean that the user has two

manufacturing times for admixtures).

CPRS A VistA computer software package called

Computerized Patient Record Systems. CPRS is an application in VistA that allows the user to enter all necessary orders for a patient in different packages from a single application. All pending orders that appear in the Unit Dose and IV modules are initially

entered through the CPRS package.

Critical Drug-Drug Interaction One of two types of drug-drug interactions identified by

order checks. The other type is a "significant" drug-

drug interaction

Cumulative Doses The number of IV doses actually administered, which

equals the total number of bags dispensed less any

Recycled, Destroyed, or Cancelled bags.

Default Answer The most common answer, predefined by the system to

save time and keystrokes for the user. The default answer appears before the two slash marks (//) and can

be selected by the user by pressing **Enter**>.

Dispense DrugThe Dispense Drug name has the strength attached to it

(e.g., Acetaminophen 325 mg). The name alone without

a strength attached is the Orderable Item name.

Delivery Times The time(s) when IV orders are delivered to the wards.

Dosage Ordered After the user has selected the drug during order entry,

the dosage ordered prompt is displayed.

DRUG ELECTROLYTES File File #50.4. This file contains the names of

anions/cations, and their concentration units.

DRUG File File #50. This file holds the information related to each

drug that can be used to fill a prescription.

Local Possible Dosages

Free text dosages that are associated with drugs that do not meet all of the criteria for Possible Dosages.

LVP

Large Volume Parenteral — Admixture. A solution intended for continuous parenteral infusion, administered as a vehicle for additive(s) or for the pharmacological effect of the solution itself. It is comprised of any number of additives, including zero, in one solution. An LVP runs continuously, with another bag hung when one bottle or bag is empty.

Manufacturing Times

The time(s) that designate(s) the general time when the manufacturing list will be run and IV orders prepared. This field in the *SIte Parameters (IV)* option (IV ROOM file, (#59.5)) is for documentation only and does not affect IV processing.

MEDICATION ADMINISTERING TEAM file

File #57.7. This file contains wards, the teams used in the administration of medication to that ward, and the rooms/beds assigned to that team.

MEDICATION INSTRUCTION file

File #51. This file is used by Unit Dose and Outpatient Pharmacy. It contains the medication instruction name, expansion, and intended use.

MEDICATION ROUTES file

File #51.2. This file contains medication route names. The user can enter an abbreviation for each route to be used at their site. The abbreviation will most likely be the Latin abbreviation for the term.

Medication Routes/ Abbreviations

Route by which medication is administered (e.g., oral). The MEDICATION ROUTES file (#51.2) contains the routes and abbreviations, which are selected by each VAMC. The abbreviation cannot be longer than five characters to fit on labels and the MAR. The user can add new routes and abbreviations as appropriate.

Non-Formulary Drugs

The medications that are defined as commercially available drug products not included in the VA National Formulary.

Non-VA Meds

Term that encompasses any Over-the-Counter (OTC) medications, Herbal supplements, Veterans Health Administration (VHA) prescribed medications but purchased by the patient at an outside pharmacy, and medications prescribed by providers outside VHA. All Non-VA Meds must be documented in patients' medical records.

Non-Verified Orders

Any order that has been entered in the Unit Dose or IV module that has not been verified (made active) by a nurse and/or pharmacist. Ward staff may not verify a non-verified order.

Orderable Item

An Orderable Item name has no strength attached to it (e.g., Acetaminophen). The name with a strength attached to it is the Dispense Drug name (e.g., Acetaminophen 325mg).

Order Check

Order checks (drug-allergy/ADR interactions, drug-drug, duplicate drug, and duplicate drug class) are performed when a new medication order is placed through either the CPRS or Inpatient Medications applications. They are also performed when medication orders are renewed, when Orderable Items are edited, or during the finishing process in Inpatient Medications. This functionality will ensure the user is alerted to possible adverse drug reactions and will reduce the possibility of a medication error.

Order Sets

An Order Set is a set of N pre-written orders. (N indicates the number of orders in an Order Set is variable.) Order Sets are used to expedite order entry for drugs that are dispensed to all patients in certain medical practices and procedures.

Order View

Computer option that allows the user to view detailed information related to one specific order of a patient. The order view provides basic patient information and identification of the order variables.

Parenteral

Introduced by means other than by way of the digestive track.

Patient Profile

A listing of a patient's active and non-active Unit Dose and IV orders. The patient profile also includes basic patient information, including the patient's name, social security number, date of birth, diagnosis, ward location, date of admission, reactions, and any pertinent remarks.

PECS

Pharmacy Enterprise Customization System. A Graphical User Interface (GUI) web-based application used to research, update, maintain, and report VA customizations of the commercial-off-the-shelf (COTS) vendor database used to perform Pharmacy order checks such as drug-drug interactions, duplicate therapy, and dosing.

Pending Order

A pending order is one that has been entered by a provider through CPRS without Pharmacy or Nursing finishing the order. Once Pharmacy or Nursing has finished and verified the order, it will become active.

PEPS

Pharmacy Enterprise Product System. A re-engineering of pharmacy data and its management practices developed to use a commercial off-the-shelf (COTS) drug database, currently First DataBank (FDB) Drug Information Framework (DIF), to provide the latest identification and safety information on medications.

Pharmacist Intervention

A recommendation provided by a pharmacist through the Inpatient Medications system's Intervention process acknowledging the existence of a critical drug-drug interaction and/or allergy/ADR interaction, and providing justification for its existence. There are two ways an intervention can be created, either via the Intervention Menu, or in response to Order Checks.

PHARMACY SYSTEM file

File # 59.7. This file contains data that pertains to the entire Pharmacy system of a medical center, and not to any one site or division.

Piggyback

Small volume parenteral solution for intermittent infusion. A piggyback is comprised of any number of additives, including zero, and one solution; the mixture is made in a small bag. The piggyback is given on a schedule (e.g., Q6H). Once the medication flows in, the piggyback is removed; another is not hung until the administration schedule calls for it.

Possible Dosages

Dosages that have a numeric dosage and numeric dispense units per dose appropriate for administration. For a drug to have possible dosages, it must be a single ingredient product that is matched to the VA PRODUCT file (#50.68). The VA PRODUCT file (#50.68) entry must have a numeric strength and the dosage form/unit combination must be such that a numeric strength combined with the unit can be an appropriate dosage selection.

Pre-Exchange Units

The number of actual units required for this order until the next cart exchange.

Primary Solution

A solution, usually an LVP, administered as a vehicle for additive(s) or for the pharmacological effect of the solution itself. Infusion is generally continuous. An LVP or piggyback has only one solution (primary solution). A hyperal can have one or more solutions.

Print Name

Drug generic name as it is to appear on pertinent IV output, such as labels and reports. Volume or Strength is not part of the print name.

Print Name{2}

Field used to record the additives contained in a commercially purchased premixed solution.

Profile

The patient profile shows a patient's orders. The Long profile includes all the patient's orders, sorted by status: active, non-verified, pending, and non-active. The Short profile will exclude the patient's discontinued and expired orders.

Prompt

A point at which the system questions the user and waits for a response.

Provider

Another term for the physician/clinician involved in the prescription of an IV or Unit Dose order for a patient.

Provider Override Reason

A reason supplied by a provider through the CPRS system, acknowledging a critical drug-drug interaction and/or allergy/ADR interaction and providing justification for its existence.

PSJI MGR

The name of the *key* that allows access to the supervisor functions necessary to run the IV medications software. Usually given to the Inpatient package coordinator.

PSJI PHARM TECH The name of the *key* that must be assigned to pharmacy

technicians using the IV module. This key allows the technician to finish IV orders, but not verify them.

PSJI PURGE The key that must be assigned to individuals allowed to

purge expired IV orders. This person will most likely be

the IV application coordinator.

PSJI RNFINISH The name of the *key* that is given to a user to allow the

finishing of IV orders. This user must also be a holder

of the PSJ RNURSE key.

PSJI USR1 The primary menu option that may be assigned to

nurses.

PSJI USR2 The primary menu option that may be assigned to

technicians.

PSJU MGR The name of the *primary menu option* and of the *key*

that must be assigned to the pharmacy package coordinators and supervisors using the Unit Dose

Medications module.

PSJU PL The name of the *key* that must be assigned to anyone

using the *Pick List Menu* options.

PSJ PHARM TECH The name of the *key* that must be assigned to pharmacy

technicians using the Unit Dose Medications module.

PSJ RNFINISH The name of the *key* that is given to a user to allow the

finishing of a Unit Dose order. This user must also be a

holder of the PSJ RNURSE key.

PSJ RNURSE The name of the *kev* that must be assigned to nurses

using the Unit Dose Medications module.

PSJ RPHARM The name of the *key* that must be assigned to a

pharmacist to use the Unit Dose Medications module. If the package coordinator is also a pharmacist he/she

must also be given this key.

Ouick Code An abbreviated for

An abbreviated form of the drug generic name (from one to ten characters) for IV orders. One of the three drug fields on which lookup is done to locate a drug. Print name and synonym are the other two. Use of

quick codes will speed up order entry, etc.

Report Device The device, identified by the user, on which computer-

generated reports selected by the user will be printed.

Schedule The frequency of administration of a medication (e.g.,

QID, QDAILY, QAM, STAT, Q4H).

Schedule Type Codes include: **O** - one time (i.e., STAT - only once),

P - PRN (as needed; no set administration times). C-continuous (given continuously for the life of the order; usually with set administration times). **R** - fill on request (used for items that are not automatically put in the cart - but are filled on the nurse's request. These can be multidose items (e.g., eye wash, kept for use by one patient and is filled on request when the supply is exhausted). And **OC** - on call (one time with no

specific time to be given, e.g., 1/2 hour before surgery).

Scheduled IV Order Inpatient Medications IV order having an

administration schedule. This includes the following IV

Types: IV Piggyback, Intermittent Syringe, IV Piggyback Chemotherapy, and Intermittent Syringe

Chemotherapy.

Self MedMedication that is to be administered by the patient to

himself.

Standard Schedule Standard medication administration schedules stored in

the ADMINISTRATION SCHEDULE file (#51.1).

Start Date/Time The date and time an order is to begin.

Status A - active, E - expired, R - renewed (or reinstated), D -

discontinued, **H** - on hold, **I** - incomplete, or **N** - nonverified, **U** - unreleased, **P** - pending, **O** - on call, **DE** - discontinued edit, **RE** - reinstated, **DR** - discontinued

renewal.

Stop Date/Time The date and time an order is to expire.

Stop Order Notices A list of patient medications that are about to expire and

may require action.

Syringe Type of IV that uses a syringe rather than a bottle or

bag. The method of infusion for a syringe-type IV may

be continuous or intermittent.

Syringe Size The syringe size is the capacity or volume of a

particular syringe. The size of a syringe is usually measured in number of cubic centimeters (ccs).

TPN Total Parenteral Nutrition. The intravenous

administration of the total nutrient requirements of the patient. The term TPN is also used to mean the solution

compounded to provide those requirements.

Units per Dose The number of Units (tablets, capsules, etc.) to be

dispensed as a Dose for an order. Fractional numbers

will be accepted.

VA Drug Class Code A drug classification system used by VA that separates

drugs into different categories based upon their

characteristics. IV cost reports can be run for VA Drug

Class Codes.

VDL Virtual Due List. This is a Graphical User Interface

(GUI) application used by the nurses when

administering medications.

Ward Group A ward group indicates inpatient nursing units (wards)

that have been defined as a group within Inpatient Medications to facilitate processing of orders.

WARD GROUP file File #57.5. This file contains the name of the ward

group, and the wards included in that group. The grouping is necessary for the pick list to be run for

specific carts and ward groups.

Ward Group Name A field in the WARD GROUP file (#57.5) used to

assign an arbitrary name to a group of wards for the

pick list and medication cart.

WARD LOCATION file File #42. This file contains all of the facility ward

locations and their related data, e.g., Operating beds, Bedsection, etc. The wards are created/edited using the

Ward Definition option of the ADT module.

(This page included for two-sided copying.)

3. Index

1

14 Day MAR, 168, 169, 170 14 Day MAR Report Example, 170

2

24 Hour MAR, 155, 156, 163 24 Hour MAR Report Example, 157

7

7 Day MAR, 162, 163, 164 7 Day MAR Report Example, 164

Α

Abbreviated Order Entry, 19, 20
Action Area, 6, 12, 18, 17, 18, 40, 65, 66, 85, 86
Action Profile #1, 174, 176
Action Profile #1 Report Example, 175
Action Profile #2, 176
Action Profile #2 Report Example, 177
Active Order List (IV), 201
Active Order List (IV) Example, 201
Activity Log, 40, 43, 47, 55, 59, 62a, 62b, 87, 98, 104, 113, 118, 119, 121b, 144, 153, 205
Activity Log Example, 55, 104, 105, 106, 150, 151
Activity Ruler, 74, 224

Additive, 68, 69, 93, 98, 111, 112, 115, 116, 117, 118, 119, 123, 146, 186, 202, 215, 217, 225, 229, 231, 233

Administration Schedule, 24, 44, 70, 94

Administration Team, 61, 121, 155, 162, 168, 174, 184, 186, 188, 202

Administration Times, 25, 24, 43, 49, 50, 51, 58, 70, 71, 89, 99, 100, 101, 110, 111, 119, 130, 140, 143, 155, 162, 168, 188, 225, 228, 230, 235

Admixture, 68, 69, 112, 114, 116, 117, 118, 143, 146, 201, 225, 226, 228, 231

Adverse Reaction Tracking (ART) Package, 29, 75

Align Labels (IV), 193

Align Labels (IV) Example, 193

Align Labels (Unit Dose), 192

Align Labels (Unit Dose) Example, 192

Allergy Indicator, 5

Allergy/ADR Example Order Checks, 124i

Asterisk, 39, 42, 43, 74, 85, 89, 144, 188

ATC, 131, 134, 136, 140

Authorized Absence, 179

Authorized Absence/Discharge Summary, 179

Authorized Absence/Discharge Summary Report Example, 179

Auto-Verify, 44, 94, 97

В

Bar Code ID, 148, 152, 193, 197, 199 Barcode ID – Return and Destroy (IV), 152 Barcode ID – Return and Destroy (IV) Example, 152 BCMA, 1, 22, 23, 40, 44, 71, 85, 94, 151, 194, 199 BCMA Units Per Dose, 22 BCMA Virtual Due List (VDL), 44, 94

C

```
Change Report/Label Devices (IV), 126
Change to Another IV Room (IV), 127
Chemotherapy, 68, 117, 118, 143, 146, 201, 226, 228
Clinic, 13, 14, 153, 155, 162, 168, 186, 192, 202
CLINIC DEFINITION file, 114, 115, 227
Clinic Group, 13, 14, 153, 155, 162, 168, 176, 186, 192, 202, 227
Clinic Location, 72
Complex Orders, 52
   Active Complex Order, 37
   Non-Verified Complex Order, 36, 81, 82
   Pending Complex Order, 36, 81, 82
Controlled Substance, 130, 164, 169
Coverage Times, 74
CPRS, 1, 13, 25, 23, 28, 39, 44, 47, 56, 58, 59, 71, 72, 85, 94, 98, 106, 111, 113, 123, 225, 228, 233
CPRS Med Order, 24, 71
CPRS Order Checks - How They Work, 219
CPRS Provider Overrides, 23
Critical Drug-Drug Interaction, 229
CWAD Indicator, 6
 D
Default Start Date Calculation, 24, 71, 111, 114, 115, 116
   Default Start Date Calculation = CLOSEST, 24, 71, 114, 115, 116
   Default Start Date Calculation = NEXT, 24, 71, 114, 115, 116
   Default Start Date Calculation = NOW, 19
   Default Start Date Calculation = NOW, 24
   Default Start Date Calculation = NOW, 68
   Default Start Date Calculation = NOW, 68
   Default Start Date Calculation = NOW, 71
   Default Start Date Calculation = NOW, 71
   Default Start Date Calculation = NOW, 114
   Default Start Date Calculation = NOW, 115
   Default Start Date Calculation = NOW, 116
   Default Start Date Calculation = NOW, 176
Default Start Date/Time, 71
Default Stop Date, 18, 19, 67, 71, 126
Default Stop Date/Time, 71
Delete Labels from Suspense (IV), 208
Delete Labels from Suspense (IV) Example, 208
Detailed Allergy/ADR List, 29, 75, 223
Discharge, 179
Discontinue All of a Patient's Orders, 60
Discontinue an Order, 12, 17, 20, 39, 40, 42, 59, 60, 64, 65, 81, 85, 87, 89, 113, 114, 118, 119, 131, 137, 144, 174, 179, 224, 234,
Discontinue an Order Example, 40, 41, 87
Discontinuing a Pending Renewal, 104
Discontinuing Duplicate Inpatient Orders, 124h
Dispense Drug, 19, 20, 21, 22, 23, 43, 44, 46, 47, 58, 69, 93, 95, 98, 111, 123, 130, 192a, 192b, 228, 232
Dispense Drug Look-Up, 215
Dispense Drug Look-Up Example, 216
Dispense Log, 55
Dispense Units Per Dose, 22
DONE Order, 24, 71, 96, 97
Dosage Ordered, 19, 21, 22, 23, 24, 25, 43, 130, 228
Drug File, 19, 58, 111, 215
Drug Inquiry (IV), 217
Drug Inquiry (IV) With Information Example, 217
Drug Inquiry (IV) With No Information Example, 217
Drug Name, 130
```

Drug Prompt, 3, 19, 20, 68, 215 Drug Text Indicator, 20, 69

Ε

Edit an Order, 42, 89 Edit an Order and Create a New Order Example, 91 Edit an Order Example, 42, 43, 89, 90 Edit Inpatient User Parameters, 125 Edit Patient's Default Stop Date, 126 Enter Units Dispensed, 134 Enter Units Dispensed Report Example, 135 Enter/Edit Allergy/ADR Data, 29, 75 Entering Barcode ID for Returns and Destroyed Medications Example, 152 Entering Returns and Destroyed Medications Example, 149 Error Information, 222 Error Messages, 221 Exiting the Order Process, 124i Expected First Dose, 58, 111 Extra Units Dispensed, 136 Extra Units Dispensed Report, 184 Extra Units Dispensed Report Example, 136, 184

F

Finish an Order, 44, 56, 58, 94, 106, 111
Finish an Order Example, 57
Finish an Order With a Duration Example, 109
Finish an Order Without a Duration Example, 107
Flag an Order Example, 59, 113
Free Text Dosage, 185
Free Text Dosage Report, 185
Free Text Dosage Report Example, 185, 186

G

Glossary, 223

Н

Header Area, 6
Hidden Actions, 4, 7, 8
History Log, 42, 55, 89, 104
Hold, 3, 12, 17, 29, 47, 48, 60, 61, 64, 65, 98, 106, 118, 130, 134, 144, 179, 224, 236
Hold All of a Patient's Orders, 60
Hold All of a Patient's Orders Example, 60
Hold an Order, 47
Hold an Order Example, 47, 48, 98
Take All of a Patient's Orders Off of Hold Example, 61
Hyperal, 68, 69, 115, 116, 117, 118, 143, 146, 201, 227, 228, 229, 233

I

Individual Labels (IV), 194
Individual Labels (IV) Example, 194, 196, 196b
Individual Order Suspension (IV), 209
Individual Order Suspension (IV) Example, 209
Infusion Rate, 69, 70, 112, 118, 119, 229
Inpatient Duplicate Therapy, 124g
Inpatient Medication Orders for Outpatients, 114, 115, 116, 155, 162, 168, 174, 176, 201, 207
Inpatient Order Entry, 11, 12, 17, 19, 20, 58, 63, 64, 65, 68, 81, 111

```
Inpatient Order Entry Example, 17, 65
Inpatient Profile, 61, 120a, 192
Inpatient Profile Example, 62b, 122
Inpatient Stop Order Notices, 176, 186, 202
Inpatient Stop Order Notices Example, 187, 203
Inpatient User Parameters, 19, 36, 44, 60, 82, 94
Inpatient Ward Parameters, 25, 24, 28, 71, 114, 115, 116, 130
Inquiries Menu, 215
Inquiries Menu Example, 215
Inquiries Option, 215
Intermittent Syringe, 70, 117
Internal Order Number, 148
Intervention, 29, 75, 225
Intervention Menu, 29, 75, 223
   Delete an Intervention Example, 32, 78
   Edit an Intervention Example, 31, 77
   New Intervention Example, 30, 76
   Print an Intervention Example, 19, 80
   View an Intervention Example, 33, 79
Introduction, 1
IRMS, 68
IV Additives, 68, 71, 114, 115, 116, 217, 230
IV Bag, 8, 148, 229
IV Drug Formulary Report (IV), 204
IV Drug Formulary Report (IV) Example, 204
IV Duration, 230
IV Flag, 58, 111
IV Fluid Orders, 112
IV Label Example, 148
IV Label Menu Example, 193
IV Menu, 63, 64, 126, 143, 193, 217
IV Menu Example, 63
IV Order
   Continuous Type, 71
   Intermittent Type, 71
IV Room, 17, 61, 65, 71, 75, 119, 121, 127, 144, 146, 148, 209, 211, 224, 230
IV Solution, 69, 217, 225
IV Stats File, 197
IV Type, 68, 69, 71, 114
 L
Label Log, 104, 119, 205
Label Menu (IV), 193
Label Print/Reprint, 192
Labels from Suspense (IV), 210
Labels from Suspense (IV) Example, 210
Large Volume Parenteral (LVP), 68, 114, 231
List Area, 6
List Manager, 5, 6, 7, 18, 40, 66, 86
Local Possible Dosages, 21, 22, 23, 24, 25, 231
Local Possible Dosages Example, 21
 М
Maintenance Options, 125
```

Maintenance Options, 125
Maintenance Options - IV, 126
Maintenance Options - Unit Dose, 125
Manufacturing List, 143, 146, 197, 211
Manufacturing List (IV), 146
Manufacturing List Example, 147
Manufacturing Record for Suspense (IV), 211

Manufacturing Record for Suspense (IV) Example, 211
Medication Administration Records (MARs), 1
Medication Routes, 23, 58, 111, 119, 130, 231
Medications Due Worksheet, 188
Medications Due Worksheet Example, 188
Menu Tree
IV Menu Tree, x
Unit Dose Menu Tree, ix
Message Window, 6

Ν

Nature of Order, 20, 28, 72 New Order Entry, 19, 68 New IV Order Entry Example, 73 New Unit Dose Order Entry Example, 27, 56 Non-Formulary Status, 21, 43, 44, 47, 58, 69, 93, 95, 98, 111 Non-Standard Schedules, 126 Non-Verified/Pending Orders, 11, 13, 18, 19, 20, 61, 67, 81 Non-Verified/Pending Orders Example, 13 NUMBER OF DAYS UNTIL STOP, 114, 115, 116

0

OCXCACHE, 219 On Call, 113 Order Actions, 40, 86 Order check, 234 data caching, 219 OCXCACHE, 219 XTMP, 219 Order Check, 123, 124 Drug-Allergy Interactions, 123, 124 Drug-Drug Interactions, 123 Duplicate Class, 123 Duplicate Drug, 123 Order Check Data Caching, 219 Order Checks, 19, 123 Drug-Allergy Interactions, 19 Drug-Drug Interactions, 19 Duplicate Class, 19 Order Checks/Interventions (OCI) Indicator, 23 Order Entry, 3, 6, 7, 11, 12, 19, 61, 63, 64 Order Lock, 11, 64 Order Options, 11 Order Set, 19, 20, 232 Orderable Item, 19, 20, 21, 23, 24, 25, 43, 44, 46, 47, 58, 69, 70, 93, 95, 98, 111, 126, 130, 186, 192a, 202, 228, 232 Orientation, 3 Other Print Info, 71

Р

Parenteral, 68, 114, 115, 225, 231, 233
Patient Action, 12, 18, 17, 18, 65, 66
Patient Information, 6, 12, 17, 37, 64, 66, 83, 223
Patient Information Example, 12, 37, 65, 66, 83, 84
Patient Lock, 11, 19, 64, 68
Patient Profile (Extended), 190
Patient Profile (Extended) Report Example, 190
Patient Profile (Unit Dose), 117, 153
Patient Profile (Unit Dose) Example, 153

Patient Profile Report (IV), 117, 205 Patient Profile Report (IV) Example, 205 Patient Profiles, 11, 63 Patient Record Update, 18, 67 Patient Record Update Example, 18, 67 Patients on Specific Drug(s) Report Example, 192a, 192b Pharmacist Intervention, 232 Pick List, 1, 44, 94, 129, 130, 131, 134, 135, 136, 138, 140, 141, 237 Pick List Menu, 3, 129 Pick List Menu Example, 129 Pick List Report, 129 Pick List Report Example, 131 Piggyback, 68, 69, 70, 115, 117, 118, 143, 146, 201, 226, 228, 233 Possible Dosages, 21, 22, 231, 233 Possible Dosages Example, 21 Pre-Exchange Units Report, 46 Production Options, 143 Profile (IV), 64, 117 Profile Report Example, 119, 120, 207 Priority 6, 14, 15a, 36a Provider, 20, 28, 112, 234 Provider Comments, 23, 24, 71 Provider Override Reason, 233 PSJ RNFINISH Key, 59, 113 PSJ RPHARM Key, 18, 29, 75, 106 PSJI PHARM TECH Key, 106 PSJU PL Key, 3, 126, 129

Q

Quick Code, 68, 116, 117, 215, 217, 230, 235

R

Regular Order Entry, 19, 20 Renew an Order, 49, 99 Active Orders, 49, 99 Complex Orders, 52 Discontinued Orders, 50, 100 Expired Continuous IV Orders, 51, 101 Expired Scheduled IV Orders, 51, 101 Expired Unit Dose Orders, 50, 100 Viewing Renewed Orders, 54 Viewing Renewed Orders, 53 Viewing Renewed Orders, 103 with CPRS Overrides/Pharmacist Interventions, 50 Renewal List (IV), 207 Renewal List (IV) Example, 207 Report Returns, 137 Reporting Medication Returns Example, 137 Reports (IV), 200 Reports (IV) Example, 200 Reports Menu, 153, 154 Reports Menu Example, 154 Reprint Labels from Suspense (IV), 210, 212 Reprint Labels from Suspense (IV) Example, 212 Reprint Pick List, 138 Reprint Pick List Example, 138 Reprint Scheduled Labels (IV), 199 Requested Start Date/Time, 57, 58, 108, 111 Returns and Destroyed Entry (IV), 148 Revision History, i

S

Sample Drug/Drug Interactions, 124o Sample Therapeutic Order Check Displays, 124s Schedule, 24, 25, 70, 118, 119, 130, 155, 162, 168, 235 Schedule Type, 25, 81, 130, 134, 186, 202, 225, 235 Scheduled Labels (IV), 197, 199 Scheduled Labels (IV) Example, 197 Screen Title, 5, 6 Select Action, 6, 7, 12, 18, 17, 56, 65, 66 Select Allergy, 29, 75 Select Order, 35b, 37, 81, 84, 223 Select Order Example, 38, 39, 84, 85 Self Med, 28 Send Pick List To ATC, 140 Service Connection, 179 Solution, 68, 69, 93, 98, 111, 112, 114, 115, 117, 119, 123, 146, 186, 202, 215, 217, 225, 229, 230, 231, 233, 234, 236 Special Instructions, 23, 24, 186, 202 Speed Actions, 59, 114 Speed Discontinue, 59, 114, 224 Speed Finish, 56, 59, 224 Speed Renew, 59, 224 Speed Verify, 59, 224 Standard Schedule, 216, 236 Standard Schedule Example, 216 Start Date/Time, 24, 40, 43, 71, 87, 114, 115, 116, 129, 130, 131, 192, 236 Stop Date/Time, 18, 24, 40, 42, 43, 55, 56, 67, 71, 72, 87, 89, 114, 115, 116, 126, 129, 130, 236 Strength, 23, 69, 118, 119, 146, 186, 202, 217, 228, 232, 233 Suspense Functions (IV), 208 Suspense Functions (IV) Menu Example, 208 Suspense List (IV), 213 Suspense List (IV) Example, 213 Syringe, 68, 93, 117, 118, 143, 146, 201, 226, 227, 228, 229, 236 Syringe Size, 117 Т

Table of Contents, v Team, 129 Test Control Codes (IV), 199 Test Control Codes (IV) Example, 199, 200 Three levels of error messages, 222 Topic Oriented Section, ix, x Total Parenteral Nutrition (TPN), 115, 229

U

Unit Dose Medications, 3, 11, 12, 125, 153, 215 Unit Dose Menu Example, 11 Units Dispensed, 130, 134, 135 Units Needed, 130 Units Per Dose, 22, 23, 24, 25, 130, 131, 140, 233 Update Daily Ward List (IV), 144 Update Daily Ward List Example, 145 Update Pick List, 141

VA Class, 192a VA Drug Class Code, 215 VA FORM 10-1158, 175, 178, 186, 202 VA FORM 10-2970, 163 VA FORM 10-5568d, 163 VDL, 24, 44, 71, 94 Verify a DONE Order (CPRS Med Order) Example, 96, 97 Verify an Order, 44, 94 Verify an Order Example, 45, 46, 95 View Profile, 12, 17, 20, 65, 66, 81, 223 View Profile Example, 20, 81 VISTA, 11, 19, 64, 68, 228 Volume, 68, 69, 93, 112, 115, 117, 118, 186, 188, 202, 230, 233, 236 Change the Volume of a Solution Example, 93

W

Ward, 14, 61, 121, 129, 153, 155, 174, 176, 184, 186, 188, 192, 202
Ward Group, 13, 14, 61, 121, 129, 130, 131, 140, 153, 155, 162, 168, 174, 176, 179, 184, 186, 188, 192, 202, 236, 237
Ward Group File, 237
Ward Group Sort

^OTHER, 13, 14, 15, 174, 176
Ward List, 63, 74, 143, 144, 146, 197, 209
Ward List (IV), 143, 199, 201
Ward List Report Example, 144
Ward Stock, 130, 164, 169



XTMP, 219