



## Colonoscopy Movi Prep Instructions

5 Days before Procedure	3 Days before Procedure	1 Day before Procedure <b>Drink clear liquids only</b> (see list on back page)	Morning of Procedure
<p><b>Stop taking these medications until after procedure:</b></p> <ul style="list-style-type: none"> <li>• Iron pill</li> <li>• Multivitamin with iron</li> </ul> <p>Resume after procedure as directed.</p>	<p><b>Stop taking these medications until after procedure:</b></p> <ul style="list-style-type: none"> <li>• Dicyclomine</li> <li>• Loperamide</li> <li>• Colestipol</li> </ul> <p>Resume after procedure as directed.</p>	<p><b>DIABETICS:</b> Take <b>HALF</b> (1/2) of usual dose of insulin. Take diabetes pills as usual.</p> <p>At <b>12 PM</b>, take <b>2</b> bisacodyl tablets. In provided container, mix one "A" packet and one "B" packet together and fill with water up to the "fill line". Shake well and refrigerate.</p> <p>From <b>4 PM to 5 PM</b> drink the entire container. Then, refill container with any type of clear liquid (not red or purple liquids) and drink from <b>5 PM to 6 PM</b>.</p> <p>Take the remaining "A" and "B" packets and put them in the container. Fill with water and refrigerate until morning.</p> <div style="text-align: center;">  <p><b>NO SOLID FOOD</b></p> </div>	<p><b>6</b> hours before procedure drink the remaining container over <b>1</b> hour.</p> <p>Refill container with any type of clear liquid (not red or purple liquids) and drink it over the next hour.</p> <p>You may drink clear liquids until <b>4</b> hours before your procedure, then nothing else by mouth!</p> <p>Take morning medications <b>4</b> hours prior to procedure:</p> <p style="text-align: center;"><b>DO NOT TAKE:</b></p> <ul style="list-style-type: none"> <li>• Diabetic pills</li> <li>• Insulin</li> <li>• Psyllium powder or seeds</li> </ul> <p>Resume medications after procedure as directed.</p> <div style="text-align: center;">  <p><b>NO SOLID FOOD</b></p> </div>

(Turn page to see clear liquid list)

## Drink plenty of fluids while on a clear liquid diet.

**\*\*NOTE: Do Not drink red or purple liquids \*\***

**\*\*NOTE: Do Not drink alcoholic beverages \*\***

### The following items **are** allowed on a clear liquid diet:

- Water (plain, carbonated or flavored)
- Black coffee or tea (**NO** cream or dairy)
- Fruit juices without pulp, such as apple, white cranberry or white grape (**NO** grapefruit, orange or red cranberry juices)
- Fruit-flavored beverages, such as lemonade (without pulp) or Kool-Aid
- Carbonated drinks, such as Sprite, 7Up, ginger ale, seltzer and dark sodas (cola and root beer)
- Sports drinks, such as Gatorade/Powerade
- Clear broths (with **NO** particles of dried food or seasoning)
- Hard candy, such as lifesavers
- Ice popsicles (without milk, fruit, seeds, or nuts)
- Gelatins, such as Jell-O
- **If you have DIABETES: Be sure to drink liquids with and without sugar**



U.S. Department of Veterans Affairs  
Veterans Health Administration  
Western North Carolina VA  
Health Care System