



Colonoscopy Magnesium Citrate + Colyte Prep Instruction

5 Days before Procedure	3 Days before Procedure	2 Days before Procedure	1 Day before Procedure Drink clear liquids only (See list on back page)	Morning of Procedure
<p>Stop taking these medications until after procedure:</p> <ul style="list-style-type: none"> • Iron pill • Multi-vitamin with iron <p>Resume after procedure as directed.</p>	<p>Stop taking these medications until after procedure:</p> <ul style="list-style-type: none"> • Dicyclomine • Loperamide • Colestipol <p>Resume after procedure as directed.</p>	<p>In the morning drink ONE bottle of magnesium citrate.</p> <p>If you take levothyroxine (thyroid medication), you must take it 4 hours after the magnesium citrate.</p>	<p>DIABETICS: Take HALF (½) of usual dose of insulin. Take diabetes pills as usual.</p> <p>At 12 PM, take 2 bisacodyl tablets, then mix prep solution and refrigerate.</p> <p>From 4 PM to 6 PM, drink HALF (½) of the prep solution.</p> <p>Refrigerate remaining solution.</p> <div style="text-align: center;">  <p>NO SOLID FOOD</p> </div>	<p>6 hours before procedure drink remaining prep solution over 2 hours.</p> <p>You may drink clear liquids until 4 hours before your procedure, then nothing else by mouth!</p> <p>Take morning medications 4 hours prior to procedure:</p> <p style="text-align: center;">DO NOT TAKE:</p> <ul style="list-style-type: none"> • Diabetes pills • Insulin • Injectables • Psyllium powder or Psyllium seeds <p>Resume after procedure as directed.</p> <div style="text-align: center;">  <p>NO SOLID FOOD</p> </div>

(Turn page to see clear liquid list)

Drink plenty of fluids while on a clear liquid diet.

****NOTE: Do Not drink red or purple liquids ****

****NOTE: Do Not drink alcoholic beverages ****

The following items **are** allowed on a clear liquid diet:

- Water (plain, carbonated or flavored)
- Black coffee or tea (**NO** cream or dairy)
- Fruit juices without pulp, such as apple, white cranberry or white grape (**NO** grapefruit, orange or red cranberry juices)
- Fruit-flavored beverages, such as lemonade (without pulp) or Kool-Aid
- Carbonated drinks, such as Sprite, 7Up, ginger ale, seltzer, and dark sodas (cola and root beer)
- Sports drinks, such as Gatorade/Powerade
- Clear broths (with **NO** particles of dried food or seasoning)
- Hard candy, such as lifesavers
- Ice popsicles (without milk, fruit, seeds, or nuts)
- Gelatins, such as Jell-O
- **If you have DIABETES: Be sure to drink liquids with and without sugar**

VA



U.S. Department of Veterans Affairs

Veterans Health Administration
Western North Carolina VA
Health Care System