

PREPARING FOR YOUR COLONOSCOPY with 2 Day Polyethylene Glycol (PEG) 3350/MiraLAX and Bisacodyl/Dulcolax

Questions after reading this?

Call the GI Triage Line at 720-723-6166 option 2, Monday – Friday and leave a message. An RN will call you back within 48 hours, or on Monday if you call on the weekend. For questions at night or on weekends, call 303-399-8020, select “0” and ask for the GI doctor on call.

As soon as you schedule your colonoscopy

- Please arrive 60 minutes before your appointment time. Expect to be here, from check to discharge, for up to four (4) hours.
- For all procedures involving **Sedation/Anesthesia**:
For your safety, an adult should accompany you home after procedures with sedation. The adult can drive you home or accompany you on public transportation. **No Uber/Lyft, taxi, bus or any other form of public transportation can be used if you are alone after sedation.**
There are a few options that can be arranged prior to your procedure if you do not have an adult to assist you. If there are safety concerns, your procedure will be canceled.
No adult to assist you? Call Veteran Transportation Service If living within 25 miles of RMR VAMC. Call (720)-857-5421/5420/5419 at least 7 days before procedure date to schedule a ride.

Prep Medications

- If you are missing your prep medications, please call the outpatient pharmacy at (888) 336-8262, choose option 1 – option 3 to request a refill. You will need to provide:
 - Name/Last 4 of SSN; Date of procedure; Prep medication name or Type of procedureAnd, if able to pick up at RMR VA Outpatient pharmacy window.

14 days before your colonoscopy

- If you haven't received a bottle of **Polyethylene glycol (PEG) 3350/MiraLAX** and four tablets of **Bisacodyl/Dulcolax** in the mail, call the GI Triage Line at 720-723-6166.
- The Anticoagulant Clinic should call you if you take a blood thinner such as Warfarin/Coumadin, Eliquis/Apixaban, Xarelto/Rivaroxaban or Pradaxa/Dabigatran. If they haven't by fourteen days prior to the procedure, call 720-723-6756.
- If you are on Plavix/Clopidogrel, Brilinta/Ticagrelor, Pletal/Cilostazol, Aggrenox/Aspirin with Dipyridamole or Prasugrel/Effient, you should receive a call from the GI Department at least fourteen days prior to the procedure. If you don't, please call 720-723-6166.

7 days before your colonoscopy

- STOP EATING corn, popcorn, frozen corn, seeds or nuts of any type. These foods are very difficult to clean from your colon.

- STOP taking any IRON supplements as they interfere with good bowel cleansing.
- **Buy two 32-ounce size bottles of Gatorade**, but NO purple, red or orange colored flavors.

3 days before your colonoscopy

- STOP taking any Vitamin E, fish oil, mineral oil, or flaxseed oil. These can cause bleeding.

2 days before your colonoscopy

- **No solid foods today!**
- Drink ONLY clear liquids (liquids you can see through) such as Gatorade, soft drinks, clear juices (without pulp), Kool-Aid, Jell-O, popsicles, clear broth, black coffee or tea. *Drink lots of clear liquids to stay hydrated! The prep will work much better if you are.*
- DO NOT DRINK milk or dairy products, alcohol, beer, or any clear liquid or Jell-O that is red, orange, or purple in color. These dyes are difficult for the doctor to see through.
- **At 3 pm**, swallow FOUR Dulcolax/bisacodyl tablets with at least 16 ounces of a clear liquid.
- **At 5 pm**, mix 1/2 (one half) of the bottle of PEG 3350/MiraLAX (about 119 grams) in one of the bottles of Gatorade. Shake it well to dissolve.



- Finish this solution **WITHIN 1 HOUR** by drinking an 8-ounce glass every 10 to 15 minutes until you finish the bottle.
- You may drink other clear liquids after you finish the PEG 3350/MiraLAX and Gatorade, up until midnight.

TIP You may want to buy flushable baby wipes and Vaseline to use during your prep for irritated skin.

The day before your colonoscopy

- **No solid foods today either!**
- Drink ONLY clear liquids (liquids you can see through) such as Gatorade, soft drinks, clear juices (without pulp), Kool-Aid, Jell-O, popsicles, clear broth, black coffee or tea. *Drink lots of clear liquids to stay hydrated! The prep will work much better if you are.*

- DO NOT DRINK milk or dairy products, alcohol, beer, or any clear liquid or Jell-O that is red, orange, or purple in color. These dyes are difficult for the doctor to see through.
- **At 3 pm**, swallow FOUR Dulcolax/bisacodyl tablets with at least 16 ounces of a clear liquid.
- **At 5 pm**, mix 1/2 (one half) of the bottle of PEG 3350/MiraLAX (about 119 grams) in one of the bottles of Gatorade. Shake it well to dissolve.
- You may drink cold or room-temperature Gatorade. Some patients find drinking the prep through a straw helpful.
- Finish this solution **WITHIN 1 HOUR** by drinking an 8-ounce glass every 10 to 15 minutes until you finish the bottle. PEG 3350/MiraLAX is a laxative that gives you diarrhea. It empties your colon, so the doctor will be able to see its walls and spot any abnormal growths (polyps).
- You may drink other clear liquids after you finish the PEG 3350/MiraLAX and Gatorade, up until midnight.
- *After midnight DON'T drink anything except what we instruct you to on the next page!*

TIP You may want to buy flushable baby wipes and Vaseline to use during your prep due to skin irritation.

The day of your colonoscopy

- **Do not eat any solid food today, and drink clear liquids only as instructed below!**
- **FIVE HOURS BEFORE LEAVING** for your colonoscopy, mix the other 1/2 (one half) of the bottle of PEG 3350/MiraLAX (about 119 grams) in the other bottle of Gatorade and shake it well to dissolve.
- Drink an 8-ounce glass of the solution every 10 to 15 minutes until you finish it.
- Take any morning medications you need at least FOUR HOURS BEFORE your colonoscopy, with a few sips of water to swallow any pills, tablets or capsules.
- *During the FOUR HOURS BEFORE your colonoscopy, for your own safety, there should be NOTHING IN YOUR MOUTH, not even chewing gum.*
- *If you have ANYTHING in those four hours, even WATER, your colonoscopy will be delayed or canceled.*
- Your stool should be clear and yellow, with no solid particles in it.



Do you have diabetes? Take special care while only drinking clear liquids!

- ✓ Monitor your blood sugar every 4 hours during the day before your colonoscopy.
- ✓ If you use insulin, the morning of the day BEFORE your colonoscopy take 1/2 (one half) your normal morning insulin dose.
- ✓ If you take diabetes medications by mouth, the morning of the day BEFORE your colonoscopy take them as usual.

- ✓ The evening BEFORE your colonoscopy, do NOT take any insulin or diabetes medications by mouth.
- ✓ The morning OF your colonoscopy, do NOT take any insulin or other diabetes medications.
- ✓ Taking any of the following GLP-1 Diabetic medications? if taking once daily, stop taking 2 days before procedure. If taking once weekly, stop 7 days before procedure: Dulaglutide, Exenatide, Liraglutide, Lixisenatide, Semaglutide, Tirzepatide. Stop taking Phentermine 7 days before procedure.

Why is following the directions so important?

Good prep: *Polyps are easier to see.*



Bad prep: *Polyps/cancer could be missed!*

